



Finite automata and pattern avoidance in words

Petter Brändén and Toufik Mansour

Abstract. We say that a word w on a totally ordered alphabet avoids the word v if there are no subsequences in w order-equivalent to v . In this paper we suggest a new approach to the enumeration of words on at most k letters avoiding a given pattern. By studying an automaton which for fixed k generates the words avoiding a given pattern we derive several previously known results for problems of this kind, as well as many new. In particular, we give a simple proof of the formula [21] for the exact asymptotics for the number of words on k letters of length n that avoids the pattern $12 \cdots (\ell+1)$. Moreover, we give the first combinatorial proof of the exact formula [9] for the number of words on k letters of length n avoiding a three letter permutation pattern.

Résumé. Soient v et w , deux mots sur un alphabet totalement ordonné. Le mot w évite le motif v si aucun sous-mot de w n'est équivalent (au sens de l'ordre) à v . Dans ce papier, nous suggérons une nouvelle approche pour énumérer les mots sur un alphabet d'au plus k lettres qui évitent un motif donné. En étudiant un automate qui engendre, pour un k fixé, tous les mots évitant un motif donné, nous obtenons des résultats nouveaux dans ce domaine, ainsi que d'autres déjà connus. En particulier, nous donnons une preuve simple de la formule de Regev pour une estimation asymptotique précise du nombre de mots de longueur n sur k lettres qui évitent le motif $12 \cdots (\ell+1)$. De plus, nous donnons pour la première fois une preuve combinatoire de la formule close de Burstein pour le calcul du nombre de mots de longueur n sur un alphabet à k lettres qui évitent un motif de permutation de 3 lettres.

1. Introduction

In this paper we study pattern avoidance in words. The subject of pattern avoidance in permutations has thrived in the last decades, see [31] and the references there. Only very recently Alon and Friedgut [3] studied pattern avoidance in words to achieve an upper bound on the number of permutations in S_n avoiding a given pattern. We study pattern avoidance in words by defining a finite automaton that generates the words avoiding a given pattern and use the transfer matrix method to count them. By this approach we are able to find the asymptotics, as $n \rightarrow \infty$, for the number of words on k letters of length n avoiding a pattern p , as well as exact enumeration results. In particular we re-derive Regev's [21] result on the exact asymptotics for the number of words on k letters of length n avoiding a pattern $12 \cdots (\ell+1)$, and give the first combinatorial proof of a formula for the number of words on k letters of length n avoiding the pattern 123.

Let S_n denote the set of permutations of the set $[n] := \{1, 2, \dots, n\}$. If $\sigma \in S_k$ and $\tau \in S_n$, we say that τ contains σ if there is a sequence $1 \leq t_1 < t_2 < \dots < t_k \leq n$ of integers such that for all $1 \leq i, j \leq k$ we

2000 Mathematics Subject Classification. 05A05, 05A15, 68Q45.

have $\tau(t_i) \leq \tau(t_j)$ if and only if $\sigma(i) \leq \sigma(j)$. Here σ is called a *pattern*. If τ does not contain σ we say that τ *avoids* σ . In the study of pattern avoidance the focus has been on enumerating and giving estimates to the number of elements in the set $S_n(\sigma)$, the set of permutations in S_n that avoids σ . Maybe the most interesting open problem in the field is: Does there exist a constant c such that $|S_n(\tau)| < c^n$ for all $n \geq 0$? This problem is equivalent to the seemingly stronger statement, see [4]:

Conjecture 1.1. (Stanley, Wilf) *For any pattern $\tau \in S_\ell$, the limit $\lim_{n \rightarrow \infty} |S_n(\tau)|^{\frac{1}{n}}$, exists and is finite.*

The conjecture has been verified for *layered* patterns [8], for all patterns which can be written as an increasing subsequence followed by a decreasing [3]. Very recently Marcus and Tardos [19] announced that they have a proof of Conjecture 1.1. In [3] Alon and Friedgut proved a weaker version of Conjecture 1.1, namely: For any permutation σ there exists a constant $c = c(\sigma)$ such that $|S_n(\sigma)| \leq c^{n\gamma^*(n)}$, where γ^* is an extremely slowly growing function, related to the Ackermann hierarchy. The method of proof in [3] was by considering pattern avoidance in words. This is also the theme of this paper.

Denote by $[k]^*$ the set of all finite words with letters in $[k]$. If $w = w_1w_2 \cdots w_s \in [k]^*$ and $v = v_1v_2 \cdots v_r \in [m]^*$ where $r \leq s$, we say that w *contains* the *pattern* v if there is a sequence $1 \leq t_1 < t_2 < \cdots < t_r \leq s$ such that for all $1 \leq i, j \leq r$ we have

$$w_{t_i} \leq w_{t_j} \quad \text{if and only if} \quad v_i \leq v_j.$$

If w does not contain v we say that w *avoids* v . For example, the word $w = 323122411 \in [4]^9$ avoids the pattern 132 and contains the patterns 123, 212, 213, 231, 312, and 321. If S is any set of finite words we denote the set of words in S that avoids v by $S(v)$.

The history of pattern avoidance in words is not as rich as the one in permutations. We mention the references [2, 3, 9, 10, 14, 21]. In [21] Regev gave a complete answer for the asymptotics for $||[k]^n(p_\ell)||$ when $n \rightarrow \infty$, where $p_\ell = 12 \cdots (\ell + 1)$ (see Theorem 4.3).

Theorem 1.2 (Regev). *For all $k \geq \ell$ we have*

$$|[k]^n(p_\ell)| \simeq C_{\ell,k} n^{\ell(k-\ell)} \ell^n \quad (n \rightarrow \infty),$$

where

$$C_{\ell,k}^{-1} = \ell^{\ell(k-\ell)} \prod_{i=1}^{\ell} \prod_{j=1}^{k-\ell} (i+j-1).$$

1.1. Organization of the paper. The paper is organized as follows. In Section 2 we present the relevant definitions and attain some preliminary results, and in Section 3 we use the transfer matrix method to determine the asymptotic growth for the sequence $n \mapsto |[k]^n(p)|$. In Section 4.1 we study the special features of the automaton, $\mathcal{A}(p_\ell, k)$, which generates the words with letters in $[k]$ that avoids the increasing pattern $12 \cdots (\ell + 1)$. Here we will give a simple proof of Theorem 4.3 using the transfer matrix method and give a combinatorial proof for the formula [9] for $|[k]^n(p)|$, where p is any permutation pattern of length three. We also consider the diagonal sequence $|[n]^n(123)|$ and determine its asymptotic growth and we also show that its generating function is transcendental. We conclude the paper by indicating further problems connected to the work in this paper.

2. Definitions and preliminary results

Given a word-pattern p and an integer $k > 0$ we define an equivalence relation \sim_p on $[k]^*$ as follows: $v \sim_p w$ if for every $r \in [k]^*$ the word vr avoids p if and only if wr avoids p . For example, if $p = 132$, $k \geq 4$, $v = 13$ and $w = 14$, then $v \sim_p w$, since 133 avoids p but 143 contains p . At first sight it may seem difficult to determine if $v \sim_p w$, since a priori there is an infinite number of right factors r to check. By the following lemma we have to check only a finite number words r .

Lemma 2.1. *Let p be a pattern of length ℓ and let $v, w \in [k]^*$ be any two words. Then $v \sim_p w$ if and only if for all words $r \in [k]^s$, $0 \leq s \leq \ell - 1$, we have*

$$vr \text{ avoids } p \quad \text{if and only if} \quad wr \text{ avoids } p.$$

PROOF. Define an equivalence relation \sim'_p on $[k]^*$ by: $v \sim'_p w$ if for all words $r \in [k]^s$, $0 \leq s \leq \ell$, we have vr avoids p if and only if wr avoids p . Clearly, $v \sim_p w$ implies $v \sim'_p w$. On the other hand if $v \sim'_p w$ we may assume that there is an $r \in [k]^*$ such that vr contains p and wr avoids p . There is at least one occurrence of p in vr that uses at most $\ell - 1$ letters of r . Thus there is a subsequence r' of r of length at most $\ell - 1$ such that vr' contains p and wr' avoids p , i.e., $v \not\sim'_p w$. \square

Let $\mathcal{E}(p, k)$ be the set of equivalence classes of \sim_p . By Lemma 2.1 the number e of equivalence classes is finite. We denote the equivalence class of a word w by $\langle w \rangle$.

Definition 2.2. Given an positive integer k and a pattern p we define a *finite automaton* (For a definition of a finite automaton, see [1] and references therein),

$$\mathcal{A}(p, k) = (\mathcal{E}(p, k), [k], \delta, \langle \varepsilon \rangle, \mathcal{E}(p, k) \setminus \{\langle p \rangle\}),$$

by

- (1) the *states* are, $\mathcal{E}(p, k)$, the equivalence-classes of \sim_p ,
- (2) $[k]$ is the *input alphabet*,
- (3) $\delta : \mathcal{E}(p, k) \times [k] \rightarrow \mathcal{E}(p, k)$ is the *transition function* defined by $\delta(\langle w \rangle, i) = \langle wi \rangle$, where wi is w concatenated with the letter $i \in [k]$,
- (4) $\langle \varepsilon \rangle$ is the *initial state*, where ε is the empty word,
- (5) all states but $\langle p \rangle$ are *final states*.

We will identify $\mathcal{A}(p, k)$ with the (labelled) directed graph with vertices $\mathcal{E}(p, k)$ and with a (labelled) edge \xrightarrow{i} between $\langle v \rangle$ and $\langle w \rangle$ if $vi \sim_p w$. Clearly, we may order the states as x_1, x_2, \dots, x_e so that if $i < j$ there is no path from x_j to x_i . The *transition matrix*, $T(p, k)$, of $\mathcal{A}(p, k)$ is the matrix of size $e \times e$ with non-negative integer coefficients defined by:

$$[T(p, k)]_{ij} = |\{s \in [k] : \delta(x_i, s) = x_j\}|.$$

Thus $[T(p, k)]_{ij}$ counts the number of edges between x_i and x_j , and $T(p, k)$ is triangular.

Example 2.3. If $p = 2314$ and $k = 5$, then it is easy to check (see [18]) that the states are $\langle \varepsilon \rangle$, $\langle 2 \rangle$, $\langle 3 \rangle$, $\langle 32 \rangle$, $\langle 34 \rangle$, $\langle 24 \rangle$, $\langle 23 \rangle$, $\langle 324 \rangle$, $\langle 341 \rangle$, $\langle 241 \rangle$, $\langle 234 \rangle$, $\langle 2342 \rangle$, $\langle 231 \rangle$, and $\langle 2314 \rangle$. Note that there are two edges between the states $\langle 324 \rangle$ and $\langle 241 \rangle$, namely $\langle 324 \rangle \xrightarrow{1} \langle 241 \rangle$ and $\langle 324 \rangle \xrightarrow{2} \langle 241 \rangle$. Moreover, all final states in $\mathcal{A}(2314, 5)$ have 3 loops, except $\langle 324 \rangle$ which has 2 loops.

The following simple lemma will be helpful in finding the asymptotic growth of the sequence $|\llbracket n \rrbracket^k(p)|$, for fixed k .

Lemma 2.4. *Let the automaton $\mathcal{A}(p, k)$ be given, let d be the number of distinct letters in p and suppose that $k \geq d - 1$. If $\langle v \rangle$ is any state different from $\langle p \rangle$, then the number of loops at $\langle v \rangle$ does not exceed $d - 1$. Moreover, there are exactly $d - 1$ loops at $\langle \varepsilon \rangle$.*

PROOF. Suppose that there are more than $d - 1$ loops at $\langle v \rangle$. Then the loops use at least d different labels. From these labels we can form a word w order-isomorphic to p . But then $vw \sim_p v$ which is a contradiction.

We may assume that the letters of p are $\{1, 2, \dots, d\}$. Let p_1 be the first letter of p . Then, if $i < p_1$ or $i > k - d + p_1$ we have $i \sim_p \varepsilon$. But there are $d - 1$ such i 's, which proves the lemma. \square

Although pattern avoidance in words and pattern avoidance in permutations share many common features, there are some important aspects in which they differ. For permutations there are three simple operations, f , that respect pattern-avoidance in the sense that $f(\tau)$ avoids $f(\sigma)$ if and only if τ avoids σ , namely the reversal, the complement and the inverse of a permutation. The first two operations have obvious generalizations to words, while the inverse does not. It has in fact been an open question to construct an inverse for words possessing “the right” properties. Such an inverse was recently constructed by Hohlweg and Reutenauer [13]. Unfortunately it is not possible to construct an inverse that respects pattern avoidance in words, which would imply the identity $||[k]^n(p)|| = ||[k]^n(p^{-1})||$, for all $k, n \geq 0$ and permutation patterns p . The first counter example to this is $||[5]^7(1342)|| = 67854 > 67853 = ||[5]^7(1423)||$. If $w \in [k]^n$ let the complement of w in $[k]^n$ be $w^c = (k+1-w_1)(k+1-w_2)\cdots(k+1-w_n)$. Then we have in fact that $\mathcal{A}(p, k)$ and $\mathcal{A}(p^c, k)$ are isomorphic as automata for any $p \in [k]^*$, since $v \sim_p w$ if and only if $v^c \sim_{p^c} w^c$.

Certainly w avoids p if and only if w^r avoids p^r , where r is the reversal operator and w and p are any words. However $\mathcal{A}(p, k)$ and $\mathcal{A}(p^r, k)$ are not in general isomorphic. Indeed, for $p = 2314$ and $k = 5$ we have that $|\mathcal{E}(2314, 5)| = 13$ and $|\mathcal{E}(4132, 5)| = 14$.

3. Transfer matrix method

In this section we use the transfer matrix method (see [27, Theorem 4.7.2]) to obtain information about the sequences $||[k]^n(p)||$. Given a matrix A let $(A; i, j)$ be the matrix with row i and column j deleted.

Theorem 3.1. *Let k be a positive integer, p be a pattern and e_k be the number of states in $\mathcal{A}(p, k)$. Let $T'(p, k) = (T(p, k); e_k - 1, e_k - 1)$. Then the generating function for $||[k]^n(p)||$ is*

$$\sum_{n \geq 0} ||[k]^n(p)|| x^n = \frac{\sum_{j=1}^{e_k-1} (-1)^{j+1} \det(I - xT', j, 1)}{\prod_{i=1}^{e_k-1} (1 - \lambda_i x)} = \frac{\det B(x)}{\prod_{i=1}^{e_k-1} (1 - \lambda_i x)},$$

where λ_i is the number of loops at state x_i , and $B(x)$ is the matrix obtained by replacing the first column in $I - xT'$ with a column of all ones.

PROOF. The theorem follows from the transfer matrix method, see [27, Theorem 4.7.2], since we want to count the number of paths of length n in $\mathcal{A}(p, k)$ from $\langle \varepsilon \rangle$ to any state other than $\langle p \rangle$ of length n in $\mathcal{A}(p, k)$. \square

Regev [21] computed the exact asymptotics for $||[k]^n(p_\ell)||$, where $p_\ell = 12 \cdots (\ell + 1)$ and $n \rightarrow \infty$. We will next find the exact asymptotics (up to a constant) for $||[k]^n(p)||$ for all patterns p . Given two sequences $\{a_n\}$ and $\{b_n\}$ of real numbers, we denote $a_n \simeq b_n$ if $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = 1$. A path in $\mathcal{A}(p, k)$ is called *simple* if it starts at $\langle \varepsilon \rangle$, does not use any loops, and does not end in $\langle p \rangle$.

Theorem 3.2. *Let p be any pattern with d distinct letters and let $k \geq d - 1$ be given. Then there is a constant $C > 0$ such that*

$$||[k]^n(p)|| \simeq C n^M (d - 1)^n \quad (n \rightarrow \infty),$$

where $M + 1$ is the maximum number of states with $d - 1$ loops, in a simple path.

PROOF. Let $P := x_1, x_2, \dots, x_j$ be a simple path in $\mathcal{A}(p, k)$. Moreover, let ℓ_j be the number of loops at state x_j . Then $||[k]^n(p)|| = \sum_P N(P, n)$ where

$$N(P, n) = \sum_{\alpha_1 + \dots + \alpha_j = n - j + 1} \ell_1^{\alpha_1} \ell_2^{\alpha_2} \cdots \ell_j^{\alpha_j},$$

and the sum is over all weak compositions of $n - j + 1$ into at most j parts. Now, $N(P, n)$ is equal to the coefficient to t^{n-j+1} in $(1 - \ell_1 t)^{-1} \cdots (1 - \ell_j t)^{-1}$. Let r be the number of i such that $\ell_i = d - 1$. Note that by Lemma 2.4 r is at least one. The dominant term of $(1 - \ell_1 t)^{-1} \cdots (1 - \ell_j t)^{-1}$ is (by partial fraction

decomposition) equal to $\frac{f(t)}{(1-(d-1)t)^r}$, where $f(t)$ is a polynomial of degree less than r and $f((d-1)^{-1}) \neq 0$. By well known results it follows that $N(P, n) \simeq C(P)(d-1)^n n^{r-1}$, where $C(P) > 0$ is a constant depending on P and k . Taking the greatest possible r yields the desired results. \square

When there are exactly $d-1$ loops at every state except $\langle p \rangle$ in $\mathcal{A}(p, k)$, it follows from Theorem 3.1 that $||[k]^n(p)|| = (d-1)^n Q(n)$, where Q is a polynomial in n . We have in fact:

Corollary 3.3. *Let $\mathcal{A}(p, k)$ be such that all states but $\langle p \rangle$ have exactly $d-1$ loops. Then*

$$|[k]^n(p)| = \sum_{j=0}^M a_j (d-1)^{n-j} \binom{n}{j},$$

where a_j counts the number of simple paths of length j in $\mathcal{A}(p, k)$. Moreover, if p is a pattern of length $\ell+1$ then $a_j = (k-d+1)^j$ for all $j = 0, 1, \dots, \ell$.

PROOF. The corollary follows from the proof of Theorem 3.2 since $N(P, n) = (d-1)^{n-j} \binom{n}{j}$. If p is a pattern of length $\ell+1$ then we have that $a_j = (k-d+1)^j$ where $j = 0, 1, \dots, \ell$, since $k^j = \sum_{i=0}^j a_i (d-1)^{j-i} \binom{j}{i}$ for all $j = 0, 1, \dots, \ell$. \square

As an example of Corollary 3.3 we note that if p is any pattern of length $\ell+1$ with exactly d different letters then $||[d]^n(p)|| = \sum_{j=0}^{\ell} (d-1)^{n-j} \binom{n}{j}$.

4. The increasing patterns

We will in this section investigate the properties of $\mathcal{A}(p_\ell, k)$, where $p_\ell = 12 \cdots (\ell+1)$. The following lemma describes the structure of $\mathcal{A}(p_\ell, k)$:

Lemma 4.1. *Let $k \geq \ell$ be given. For any subset S of $[k]$ of size ℓ let w_S be the word consisting of the elements of S listed in increasing order. Then the words w_S together with p_ℓ constitute a complete set of representatives for the equivalence-classes $\mathcal{E}(p_\ell, k)$. In particular we have:*

$$|\mathcal{E}(p_\ell, k)| = \binom{k}{\ell} + 1.$$

If $S = \{s_1 < \cdots < s_\ell\} \subseteq [k]$ and $j \in [k]$ let $S^j = \{s_1 < \cdots < s_{i-1} < j < s_{i+1} < \cdots < s_\ell\}$, where i is the integer such that $s_{i-1} < j \leq s_i$ ($s_0 := 0, s_{\ell+1} := k+1$). Then

$$\delta(\langle w_S \rangle, j) = \begin{cases} \langle w_{S^j} \rangle & \text{if } j \leq s_\ell, \\ \langle p_\ell \rangle & \text{otherwise.} \end{cases}$$

In particular, the loops of w_S are the elements of S .

PROOF. It is clear that the words w_S are representatives for different classes. Let $v \in [k]^*(p_\ell)$. We say that an increasing subword $x_1 x_2 \cdots x_j$ of v is *extendible* if $x_j \leq k+j-\ell-1$, i.e., if we may extend $x_1 x_2 \cdots x_j$ to an occurrence of p_ℓ using letters from $[k]$. Suppose that the maximum length of an extendible increasing subsequence in v is equal to s , $s \leq \ell$. For $1 \leq j \leq s$ let

$$r_j(v) := \min\{x_j : x_1 x_2 \cdots x_j \text{ is an extendible subword of } v\}.$$

Clearly $r_1(v) < r_2(v) < \cdots < r_s(v)$. Let

$$S = \{r_1(v), r_2(v), \dots, r_s(v), k+s+1-\ell, k+s+2-\ell, \dots, k\}.$$

Then we see that $w_S \sim v$. The statement about the transition function follows from the construction. \square

In the sequel we will use some standard notation from the theory of partitions and symmetric functions. For undefined terminology we refer the reader to Chapter 7 of [28].

Theorem 4.2. *Define a partial order on the final states in $\mathcal{A}(p_\ell, k)$ as follows: $x \leq y$ if there exists a path from x to y in $\mathcal{A}(p_\ell, k)$. Then this partial order is isomorphic to $J([\ell] \times [k - \ell])$, the lattice of order ideals of the poset $[\ell] \times [k - \ell]$.*

PROOF. Let $S = \{s_1 < s_2 < \dots < s_\ell\}$ and $T = \{t_1 < t_2 < \dots < t_\ell\}$ be subsets of $[k]$. We claim that there exists a path from $\langle w_S \rangle$ to $\langle w_T \rangle$ if and only if $s_i \geq t_i$ for all $1 \leq i \leq \ell$. From this the theorem follows since the latter poset is isomorphic to the interval $[\emptyset, \lambda_{\ell, k-\ell}]$, in the Young's lattice, where $\lambda_{\ell, k-\ell} := (k - \ell, k - \ell, \dots, k - \ell)$ is of length ℓ . Indeed, consider the bijection defined by:

$$(s_1, s_2, \dots, s_\ell) \mapsto (s_\ell - \ell, s_{\ell-1} - \ell + 1, \dots, s_1 - 1) \in [\emptyset, \lambda_{\ell, k-\ell}].$$

Then $s_i \geq t_i$ for all $1 \leq i \leq j$ if and only if the image of S is greater than the image of T in $[\emptyset, \lambda_{\ell, k-\ell}]$. But $[\emptyset, \lambda_{\ell, k-\ell}]$ is its own dual, so the statement follows from the simple fact that $[\emptyset, \lambda_{\ell, k-\ell}]$ is isomorphic to $J([\ell] \times [k - \ell])$.

If there is an edge between $\langle w_S \rangle$ and $\langle w_T \rangle$, we are done by Lemma 4.1. The “only if” direction thus follows by induction on the length of the path.

Now, if $s_i \geq t_i$ for all $1 \leq i \leq \ell$ consider the path

$$\langle w_S \rangle \xrightarrow{t_1} \langle w_{St_1} \rangle \xrightarrow{t_2} \langle w_{St_1 t_2} \rangle \xrightarrow{t_3} \dots \xrightarrow{t_\ell} \langle w_{St_1 t_2 \dots t_\ell} \rangle.$$

It is not hard to see that $\langle w_{St_1 t_2 \dots t_\ell} \rangle = \langle w_T \rangle$, which completes the proof. \square

We now have a different proof of the following theorem of Regev [21]:

Theorem 4.3 (Regev). *Let $C_{\ell, k}^{-1} = \ell^{\ell(k-\ell)} \prod_{i=1}^{\ell} \prod_{j=1}^{k-\ell} (i+j-1)$. For all $k \geq \ell$ we have*

$$|[k]^n(p_\ell)| \simeq C_{\ell, k} n^{\ell(k-\ell)} \ell^n \quad (n \rightarrow \infty).$$

PROOF. By Corollary 3.3 and Theorem 4.2 we have that

$$|[k]^n(p_\ell)| \simeq a_M \ell^{-M} \binom{n}{M} \ell^n \simeq \frac{a_M}{M!} \ell^{-M} n^M \ell^n \quad (n \rightarrow \infty),$$

where $M = \ell(k - \ell)$ and a_M is equal to the number of maximal chains in $J([\ell] \times [k - \ell])$. By [28, Proposition 7.10.3] and the hook-length formula [28, Corollary 7.21.6] we have that

$$a_{\ell(k-\ell)} = f^{\lambda_{\ell, k-\ell}} = \frac{(\ell(k-\ell))!}{\prod_{i=1}^{\ell} \prod_{j=1}^{k-\ell} (i+j-1)},$$

from which the theorem follows. \square

It should be clear from the correspondence in Theorem 4.2 that the simple paths of length r in $\mathcal{A}(p_\ell, k + \ell)$ are in a one-to-one correspondence with tableaux T of the following type:

- (i) T is weakly increasing in rows and columns,
- (ii) no integer appears in more than one row,
- (iii) the entries of T are exactly $[r]$,
- (iv) the shape of T is contained in $\lambda_{\ell, k}$.

Recall that the tableaux satisfying (i) and (ii) above are the *border-strip* tableaux (or *rim-hook* tableaux)

of height zero. We call these tableaux *segmented*. Let $a(\ell, k, r)$ denote the number of segmented tableaux satisfying (iii) and (iv), so that:

$$|[k + \ell]^n(p_\ell)| = \sum_{r=0}^{\ell k} \ell^{n-r} a(\ell, k, r) \binom{n}{r}. \tag{4.1}$$

The function $a(\ell, k, r)$ is actually a polynomial in k of degree r . To see this let us call a segmented tableau inside $[\ell] \times [k]$ *primitive* if all columns are different, and let the set of such tableaux of length i with r different entries be $\mathcal{PR}_{\ell, i, r}$. If we denote the number of elements in $\mathcal{PR}_{\ell, i, r}$ by $\text{pr}(\ell, i, r)$ we have

$$a(\ell, k, r) = \sum_{i=r/\ell}^r \text{pr}(\ell, i, r) \binom{k}{i},$$

since for any such primitive tableaux of length i we may insert α_1 copies of the first column before the first column, α_2 copies of the second column between the first and the second column, and so on. After the last column we may insert α_{i+1} columns of all blanks, requiring that $\alpha_1 + \alpha_2 + \dots + \alpha_{i+1} = k - i$. Thus there are $\binom{k}{i}$ segmented tableaux arising from a given primitive one. The numbers $\text{pr}(\ell, i, r)$ are in general hard to count, but there are two special cases which are nice, namely $\text{pr}(\ell, r, r)$ and $\text{pr}(2, i, r)$. We start by counting $\text{pr}(\ell, r, r)$.

Theorem 4.4. *With definitions as above: $\text{pr}(\ell, n, n) = |S_n(p_\ell)|$.*

PROOF. We will define a bijection between S_n and $\cup_{\ell \geq 0} \mathcal{PR}_{\ell, n, n}$ such that the height of the tableau corresponds to the greatest increasing subsequence in the permutation. Recall the definition of $r_i(v)$ in the proof of Lemma 4.1, and let $r(v) = (r_1(v), r_2(v), \dots, r_\ell(v))$, where ℓ is the length of the longest increasing subsequence in v . Let k be large enough so that all increasing subsequences in permutations in S_n are considered extendible.

Now, if $\pi = \pi_1 \pi_2 \dots \pi_n$ is any permutation in S_n define $T = T(\pi)$ as follows. Let the first column of T be $r(\pi)$, the second column be $r(\pi_1 \dots \pi_{n-1})$, and so on. The image of the permutation 351462 is:

$$\begin{array}{cccccc} 1 & 1 & 1 & 1 & 3 & 3 \\ T(351462) = & 2 & 4 & 4 & 5 & 5 \\ & 6 & 6 & & & \end{array}$$

By Lemma 4.1 we have that $T(\pi) \in \mathcal{PR}_{\ell, n, n}$. Moreover from Lemma 4.1 we also get that a tableau T is the image of some $\pi \in S_n$ if and only if

- (a) T has n columns and entries $1, 2, \dots, n$,
- (b) Let T^i denote the i th column. If $i < j$ then T^i is smaller than T^j in the product order. (If T^i and T^j have different size fill the empty slots of T^j with $n + 1$),
- (c) Exactly one new entry appears every time you move from T^{i+1} to T^i .

Now, if $T \in \cup_{\ell \geq 0} \mathcal{PR}_{\ell, n, n}$ condition (a) and (b) are trivially satisfied. At least one new entry appears every time we move from T^{i+1} to T^i , since otherwise $T^i = T^{i+1}$ and T fails to be primitive. On the other hand if there appears more than one new entry in a transition then in a later transition there must appear no new entry, since T has n columns and n distinct entries. This verifies condition (c) and the theorem follows. \square

A special case of Theorem 4.4 is that $\text{pr}(2, n, n) = C_n$, the n th Catalan number. This is also a special case of the next theorem. Note that Theorem 4.5 is what we need to have combinatorial proof of a closed formula, see Theorem 4.7, for the numbers $|[k]^n(123)|$. Burstein [9] achieved a different, but of course equivalent, formula for $|[k]^n(123)|$, but not in a bijective manner.

Theorem 4.5. *With definitions as above: $\text{pr}(2, i, r) = \frac{1}{i+1} \binom{2i}{i} \binom{i}{r-i}$.*

Before we give a proof of Theorem 4.5 we will need some definitions and a lemma. Let $\mathcal{PR}^+(2, s, r)$ be the tableaux in $\mathcal{PR}(2, s, r)$ that fill up the shape $[2] \times [r]$, and let $\text{pr}^+(2, s, r) := |\mathcal{PR}^+(2, s, r)|$. Then $\text{pr}(2, s, r) = \text{pr}^+(2, s, r) + \text{pr}^+(2, s, r + 1)$ since we get the tableaux that do not fill up the shape by deleting all entries $r + 1$. To prove the theorem we will show that $\text{pr}^+(2, s, r) = \binom{s-1}{2s-r} C_s$, where C_s is the s th Catalan number.

We first define an operation $+$ that takes tableaux with r different entries to tableaux with $r + 1$ different entries. Let $T \in \mathcal{PR}^+(2, s, r)$. Suppose that j is an index such that $T_{ij} = T_{i(j+1)}$ for some $i = 1, 2$. Write T as $T = LR$ where L is the j first columns and R is the $s - j$ last columns. Let R' be the array order equivalent to R with entries the same as R , add $r + 1$, take away $T_{i(j+1)}$ (two arrays A and B are said to be order equivalent if $A_{ij} \leq A_{i'j'}$ if and only if $B_{ij} \leq B_{i'j'}$ for all i, j, i', j'). We define $T + j$ to be the tableaux $T + j := LR'$. In T there are exactly $t = 2s - r$ indices $j \in [s - 1]$ such that $T_{ij} = T_{i(j+1)}$ for some $i = 1, 2$. Let $S = \{s_1 < s_2 < \dots < s_t\}$ be these indices and define a function $\Phi : \mathcal{PR}^+(2, s, r) \rightarrow \binom{[s-1]}{t} \times \mathcal{ST}_{2,s}$, where $\mathcal{ST}_{2,s}$ is the set of standard tableaux of shape $[2] \times [s]$, by

$$\Phi(T) = (S, T + s_t + s_{t-1} + \dots + s_1).$$

The fact that Φ is a bijection will prove the theorem, since by the hook-length formula we have $|\mathcal{ST}_{2,s}| = C_s$. To find the inverse of Φ we need a kind of inverse operation to $+$.

Let $T \in \mathcal{PR}^+(2, s, r)$ and $1 \leq b \leq s - 1$ be such that $T_{1b} < T_{1(b+1)}$ and $T_{2b} < T_{2(b+1)}$. Define two arrays $T|_b$ and $T|^b$ as follows. Write $T = LR$ where L are the b first columns and R are the $s - b$ last columns. Define $T|^b := L'R'$, to be the array where $L = L'$ and R' is the unique array order equivalent to R , with entries the same as R add T_{1b} take away r . Similarly, let $T|_b := L'R'$, be the array with $L = L'$ and where R' is the unique array order equivalent with R , with entries the same as R , add T_{2b} take away r .

$$\begin{array}{c} 1\ 2\ 4\ 4 \\ 3\ 5\ 6\ 7 \end{array} \Big|_2^2 = \begin{array}{c} 1\ 2\ 2\ 2 \\ 3\ 5\ 4\ 6 \end{array}$$

$$\begin{array}{c} 1\ 2\ 4\ 4 \\ 3\ 5\ 6\ 7 \end{array} \Big|_2 = \begin{array}{c} 1\ 2\ 4\ 4 \\ 3\ 5\ 5\ 6 \end{array}$$

Note that exactly one of $T|^2$ and $T|_2$ above is a primitive segmented tableaux. This is no accident.

Lemma 4.6. *Let $T \in \mathcal{PR}^+(2, s, r)$ and $1 \leq b \leq s - 1$ be such that $T_{1b} < T_{1(b+1)}$ and $T_{2b} < T_{2(b+1)}$. Then*

$$T|_b \in \mathcal{PR}^+(2, s, r - 1) \Leftrightarrow T|^b \notin \mathcal{PR}^+(2, s, r - 1) \Leftrightarrow T_{2(b+1)} = T_{2b} + 1.$$

Moreover, if $B = T|^b \in \mathcal{PR}^+(2, s, r - 1)$ then $B_{1b} = B_{1(b+1)}$ and if $A = T|_b \in \mathcal{PR}^+(2, s, r - 1)$ then $A_{1b} = A_{1(b+1)}$.

PROOF. Consider $A := T|_b$. All entries in T that are smaller than T_{2b} will be mapped onto themselves and $A_{ij} = T_{ij} - 1$ for $A_{ij} > T_{2b}$. Therefore $A \in \mathcal{PR}^+(2, s, r - 1)$ if and only if $T_{2(b+1)} = T_{2b} + 1$ (since otherwise the entry T_{2b} will appear in both the first and the second row).

Consider $B := T|^b$. Let $y_i, i = 1, 2, \dots, h$ be the entries in T satisfying $T_{2b} < y_i \leq T_{2(b+1)}$ ordered by size. Then the entry y_1 will be mapped to an element smaller than T_{2b} and y_i will be mapped to y_{i-1} for $i > 1$. Thus $B \in \mathcal{PR}^+(2, s, r - 1)$ if and only if $T_{2(b+1)} > T_{2b} + 1$ as claimed.

The last statement is a direct consequence of the above proof. \square

We are now ready to give a proof of Theorem 4.5.

PROOF OF THEOREM 4.5. If $T \in \mathcal{PR}^+(2, s, r)$ and $1 \leq b \leq s - 1$ are such that $T_{1b} < T_{1(b+1)}$ and $T_{2b} < T_{2(b+1)}$ we define $T - b$ to be the one of the arrays $T|_b$ and $T|^b$ which is in $\mathcal{PR}^+(2, s, r - 1)$. By

Lemma 4.6 we have that

$$\begin{aligned} (T + j) - j = T & \quad \text{if } T_{ij} = T_{i(j+1)} \quad \text{for some } i = 1, 2, \\ (T - j) + j = T & \quad \text{if } T_{ij} < T_{i(j+1)} \quad \text{for both } i = 1, 2. \end{aligned} \tag{4.2}$$

Now, if $S = \{x_1 < x_2 < \dots < x_t\}$, where $t = 2s - r$ and $P \in \mathcal{ST}_{2,s}$ we let

$$\Psi(S, P) := P - x_1 - x_2 - \dots - x_t.$$

By 4.2 it follows that Ψ is the inverse to Φ and the theorem follows. \square

We now have a combinatorial proof of the following theorem given in a different form in [9]:

Theorem 4.7. *For all $n, k \geq 0$ we have*

$$|[k + 2]^n(123)| = \sum_{r,i} 2^{n-r} C_i \binom{i}{r-i} \binom{n}{r} \binom{k}{i},$$

where C_i is the i th Catalan number. The generating function

$$F(x, y) := \sum_{n,k} |[k + 2]^n(123)| x^k y^n,$$

is given by

$$F(x, y) = \frac{1}{(1-x)(1-2y)} C\left(\frac{xy(1-y)}{(1-x)(1-2y)^2}\right),$$

where $C(z)$ is the generating function for the Catalan numbers. Equivalently, $F(x, y)$ is algebraic of degree two and satisfies the equation:

$$x(1-x)y(1-y)F^2 - (1-x)(1-2y)F + 1 = 0.$$

To complete the picture for permutation patterns of length 3 it remains to enumerate $|[k]^n(132)|$. Simion and Schmidt [25] introduced a simple bijection between $S_n(123)$ and $S_n(132)$ which fixes each element of $S_n(123) \cup S_n(132)$. West [30] generalized this bijection to obtain a bijection between $S_n(p)$ and $S_n(q)$ where $p(\ell) = q(\ell - 1) = \ell$, $p(\ell - 1) = q(\ell) = \ell - 1$, and $p, q \in S_\ell$. This bijection, in turn, generalizes to words as follows.

Theorem 4.8. *Let $p = p_1 p_2 \dots p_\ell$ be a pattern with greatest entry equal to d and $p_{\ell-1} = d - 1$, $p_\ell = d$. If d occurs exactly once in p then*

$$|[k]^n(p)| = |[k]^n(\tilde{p})|,$$

where $\tilde{p} = p_1 p_2 \dots p_\ell p_{\ell-1}$.

PROOF. The proof is a straight forward generalization of West's algorithm presented in [30, Sec. 3.2]. \square

For example, if $p = 132$ then $\tilde{p} = 123$. Hence, by Theorem 4.8 we get that if p and q are any permutation patterns of length 3 then $|[k]^n(p)| = |[k]^n(q)|$ for all $n, k \geq 0$ (see [9] for an analytical proof). If $p = 1232$ then $\tilde{p} = 1223$. Hence, Theorem 4.8 gives $|[k]^n(1232)| = |[k]^n(1223)|$ for all $n, k \geq 0$.

Since, $S_n(p) \subset [n]^n(p)$, the numbers $|[n]^n(p)|$ are interesting. A sequence $f(n)$ is *polynomially recursive* (P -recursive) if there is a finite number of polynomials $P_i(n)$ such that $\sum_{i=0}^N P_i(n) f(n+i) = 0$, for all integers $n \geq 0$. For the case when p is permutation pattern of length 3 we have the following:

Theorem 4.9. *Let p be a permutation pattern of length 3. Then the sequence $f(n) := |[n]^n(p)|$ is P -recursive and satisfies the three term recurrence:*

$$p(n)f(n-2) + q(n)f(n-1) + r(n)f(n) = 0,$$

where

$$\begin{aligned} p(n) &= 3(n-3)(n-1)(3n-5)(3n-4)(5n-4), \\ q(n) &= 288 - 1440n + 2780n^2 - 2435n^3 + 976n^4 - 145n^5, \quad \text{and} \\ r(n) &= 2(n-2)^2n(n+1)(5n-9). \end{aligned}$$

PROOF. The fact that $f(n)$ is P -recursive follows easily from the expansion of $f(n)$ as a double sum using Theorem 4.7 and the theory developed in [17]. The polynomials p, q and r were found using the package MULTISUM (see [29]) developed by Wegschaider and Riese. \square

Corollary 4.10. *The asymptotics of $f(n) = |[n]^n(123)|$ is given by $f(n) \sim Cn^{-2} \left(\frac{27}{2}\right)^n$, where $C > 0$ is a constant.*

PROOF. This is a direct consequence of Theorem 4.9 and the theory of asymptotics for P -recursive sequences, see [32]. \square

A consequence of this is that the generating function of $f(n)$ is transcendental, since the exponent of n in the asymptotic expansion of a sequence with an algebraic generating function is never a negative integer.

4.1. Generating function approach. In this section we will investigate the generating function that enumerates the number of segmented tableaux according to size of rows and number of different entries. Let $A_\ell(x_1, x_2, \dots, x_\ell, t)$ be the generating function:

$$A_\ell = \sum_T x_1^{\lambda_1(T)} x_2^{\lambda_1(T) - \lambda_2(T)} \dots x_\ell^{\lambda_{\ell-1}(T) - \lambda_\ell(T)} t^{N(T)},$$

where $\lambda_i(T)$ denotes the size of row i in T , $N(T)$ denotes the number of different entries in T and the sum is over all segmented tableaux with at most ℓ rows. For $i = 1, 2, \dots, \ell$ let $A_\ell^i(x_1, \dots, x_\ell, t)$ be the generating function for those tableaux which have their maximal entry in row i . If $F(x_1, x_2, \dots, x_n)$ is a formal power-series in n variables the *divided difference* of F with respect to the variable x_i is $\Delta_i F := \frac{F - F(x_i=0)}{x_i}$, where $F(x_i=0)$ is short for $F(x_1, x_2, \dots, x_{i-1}, 0, x_{i+1}, \dots, x_n)$.

Theorem 4.11. *With definitions as above we have that A_ℓ satisfies the following system of equations:*

$$\begin{aligned} A_\ell &= 1 + A_\ell^1 + \dots + A_\ell^\ell, \\ A_\ell^1 &= x_1 x_2 t A_\ell + x_1 x_2 A_\ell^1, \\ A_\ell^2 &= x_3 t \Delta_2 A_\ell + x_3 \Delta_2 A_\ell^2, \\ &\vdots \\ A_\ell^{\ell-1} &= x_\ell t \Delta_{\ell-1} A_\ell + x_\ell \Delta_{\ell-1} A_\ell^{\ell-1}, \\ A_\ell^\ell &= t \Delta_\ell A_\ell + \Delta_\ell A_\ell^\ell. \end{aligned}$$

PROOF. The theorem follows by treating two separate cases. Let n be the greatest entry in the tableau T . The case when there is one n in a row corresponds to the first summand and the case when there are more than one n in a row corresponds to the second summand. \square

When $\ell = 2$, $A = A_2$, the system boils down to:

$$\left((1 - x_2^{-1}) \left(1 - \frac{x_1 x_2 t}{1 - x_1 x_2} \right) - x_2^{-1} t \right) A = 1 - x_2^{-1} (1 + t) A(x_2 = 0). \quad (4.3)$$

This equation can be solved using the so called *kernel method* as described in [5]. If we let

$$x_2 = \frac{1 + x_1(1 + 2t) - \sqrt{(1 + x_1(1 + 2t))^2 - 4x_1(1 + t)^2}}{2x_1(1 + t)},$$

then the parenthesis in front of A in 4.3 cancels, and we get:

$$A(x_2 = 0) = \frac{1 + x_1(1 + 2t) - \sqrt{(1 + x_1(1 + 2t))^2 - 4x_1(1 + t)^2}}{2x_1(1 + t)^2}.$$

By the interpretation of $a(\ell, k, r)$, we have that the bi-variate generating function for $a(2, k, r)$ is $(1 + x_1)^{-1}A_2(x_1, 1, t)$. From this and 4.1 one may derive an analytic proof of Theorem 4.7.

5. Further results and open problems

5.1. Further directions. Recall that the Stanley-Wilf Conjecture asserts that for any permutation π the limit $\lim_{n \rightarrow \infty} |S_n(\pi)|^{1/n}$ exists and is finite. What about the sequence $|[n]^n(\pi)|$?

Problem 5.1. *Let π be a permutation. Is there a constant $0 < C < \infty$ such $|[n]^n(\pi)| \leq C^n$ for all $n \geq 0$?*

Note that the answer to Problem 5.1 is no when π is not a permutation, since then $S_n = S_n(\pi) \subseteq [n]^n(\pi)$. Again, Problem 5.1 is equivalent to the statement that

$$\lim_{n \rightarrow \infty} |[n]^n(\pi)|^{1/n},$$

exists and is finite. This is because for all $m, n \geq 0$ we have

$$|[n + m]^{n+m}(\pi)| \geq |[n]^n(\pi)| \cdot |[m]^m(\pi)|,$$

so we may apply Fekete’s Lemma on sub-additive sequences. See [4, Theorem 1] for details (the proof extends to words word for word). For permutations $\pi \in S_3$ we have by Corollary 4.10 that $\lim_{n \rightarrow \infty} |[n]^n(\pi)|^{1/n} = 27/2$ as opposed to $\lim_{n \rightarrow \infty} |S_n(\pi)|^{1/n} = 4$.

For which permutations do we know Problem 5.1 holds? It follows from the work in [3] Problem 5.1 holds for all permutations which can be written as an increasing sequence followed by a decreasing. Also, with no great effort Bóna’s proof [8] of the Stanley-Wilf conjecture for layered patterns may be extended to this setting. Thus for all classes that the Stanley-Wilf conjecture is known to hold, the seemingly stronger Problem 5.1 holds. The following conjecture therefore seems plausible:

Conjecture 5.2. *For all permutations π we have:*

$$\exists C \forall n (|[n]^n(p)| \leq C^n) \Leftrightarrow \exists D \forall n (|S_n(p)| \leq D^n).$$

There are several problems concerning the automaton associated to a pattern that has connections to the above problems. One problem is to give an estimate to the number of simple paths in $\mathcal{A}(p, k)$, another is to estimate the number of equivalence classes in $\mathcal{A}(p, k)$. Yet another problem is to give an estimate to the maximum size of an equivalence class.

5.2. Formula for $|[k]^n(p)|$. Our algorithm (see Theorem 3.1) for finding a formula for $|[k]^n(p)|$ is implemented in C++ and Maple, see [18]. The first with input p and k and output the automaton $\mathcal{A}(p, k)$ and the second with input the automaton $\mathcal{A}(p, k)$ and output the exact formula for $|[k]^n(p)|$. This algorithm allows us to get an explicit formula for $|[k]^n(p)|$ where $p \in S_k$ and $k \geq 1$ are given. For example, an output for the algorithm for $p \in S_4$ and $k = 3, 4, 5, 6$ is given by [18].

Finally we remark that our method can be generalized as follows. Given a set of patterns T we define an equivalence relation \sim_T on $[k]^*$ by: $v \sim_T w$ if for all words $r \in [k]^*$ we have vr avoids T if and only if wr avoids T , where a word u avoids T if u avoids all patterns in T . As in Section 2 we define an automaton $\mathcal{A}(T, k)$ with the equivalence classes of \sim_T as states. With minor changes in the proof, Theorem 3.1 can be extended to avoidance of a set of patterns. For example, if $T = \{1234, 2134\}$ and $k = 6$, then by [18] we get that

$$(5.1) \quad |[6]^n(T)| = 4 \cdot 3^n + 12 \binom{n}{2} 3^{n-2} + 24 \binom{n}{3} 3^{n-3} + 54 \binom{n}{4} 3^{n-4}$$

$$(5.2) \quad + 60 \binom{n}{5} 3^{n-5} + 40 \binom{n}{6} 3^{n-6} - 3 \cdot 2^n.$$

Acknowledgements. We would like to thank M. Bousquet-Mélou and C. Krattenthaler for interesting discussions, D. Zeilberger for informing us about the work of Wegschaider and Riese, and Wegschaider and Riese for running their program for us.

References

- [1] A. V. AHO, R. SETHI, J.D. ULLMAN, Compilers: principles, techniques and tools, Addison-Wesley, Reading, Mass., 1986.
- [2] M. ALBERT, R. ALDRED, M.D. ATKINSON, C. HANDLEY, AND D. HOLTON, Permutations of a multiset avoiding permutations of length 3, *Europ. J. Combin.* **22** (2001) 1021–1031.
- [3] N. ALON AND E. FRIEDGUT, On the number of permutations avoiding a given pattern, *J. Combin. Theory Series A* **89** (2000) 133–140.
- [4] R. ARRATIA, On the Stanley-Wilf conjecture for the number of permutations avoiding a given pattern, *Elect. J. Combin.* **6** (1999) (electronic)
- [5] C. BANDERIER AND M. BOUSQUET-MÉLOU AND A. DENISE, AND P. FLAJOLET AND D. GARDY AND D. GOUYOU-BEAUCHAMPS, Generating functions for generating trees, Formal power series and algebraic combinatorics (Barcelona, 1999). *Discr. Math.* **246** (2002) 29–55.
- [6] M. BÓNA, Permutations avoiding certain patterns: the case of length 4 and some generalization, *Discr. Math.* **80** (1997) 257–272.
- [7] M. BÓNA, Exact enumeration of 1342-avoiding permutations: A close link with labelled trees and planar maps, *J. Combin. Theory Series A* **175** (1997) 55–67.
- [8] M. BÓNA, The solution of a conjecture of Stanley and Wilf for all layered patterns, *J. Combin. Theory Series A* **85** (1999) 96–104.
- [9] A. BURSTEIN, Enumeration of words with forbidden patterns, Ph.D. thesis, University of Pennsylvania, 1998.
- [10] A. BURSTEIN AND T. MANSOUR, Words restricted by patterns with at most 2 distinct letters, *Elect. J. Combin.* **9:2** (2002) #R3.
- [11] I.M. GESSEL, Symmetric functions and P -recursiveness, *J. Combin. Theory Series A* **53** (1990) 257–285.
- [12] J. M. HAMMERSLEY, A few seedlings of research, *Proceedings Sixth Berkeley Symposium on Mathematical Statistics and Probability*, Volume **1**, Berkeley/Los Angeles, 1972, University of California Press, 345–394.
- [13] C. HOHLWEG AND C. REUTENAUER, Inverses of words and the parabolic structure of the symmetric group, *Europ. J. Combin.* **22** (2001) 1075–1082.
- [14] M. KLAZAR, The Füredi-Hajnal conjecture implies the Stanley-Wilf conjecture, Formal power series and algebraic combinatorics (Moscow,2000), Springer, Berlin (2000) 250–255
- [15] D.E. KNUTH, The art of computer programming, Volume **1**, Fundamental algorithms, Addison Wesley, Reading, Massachusetts, 1973.
- [16] D.E. KNUTH, The art of computer programming, Volume **3**, Fundamental algorithms, Addison Wesley, Reading, Massachusetts, 1973.
- [17] L. LIPSHITZ, D -finite power series, *J. Algebra* **122** (1989) 353–373.
- [18] T. MANSOUR, Finite automata and pattern avoidance, Maple Programming, Available in <<http://math.haifa.ac.il/toufik>>.
- [19] A. MARCUS AND G. TARDOS, On 0-1 matrices and small excluded submatrices, <<http://www.renyi.hu/~tardos/submatrix.ps>>, 2003.
- [20] A. REGEV, Asymptotic values for degrees associated with strips of Young diagrams, *Adv. Math.* **41** (1981) 115–136.
- [21] A. REGEV, Asymptotics of the number of k -words with an ℓ -descent, *Elect. J. Combin.* **5** (1998) #R15.
- [22] D. G. ROGERS, Ascending sequences in permutations, *Discr. Math.* **22** (1978) 35–40.
- [23] D. ROTEM, On a correspondence between binary trees and a certain type of permutation, *Info. Proc. Letters* **4** (1975) 58–61.
- [24] D. ROTEM, Stack sortable permutations, *Discr. Math.* **33** (1981) 185–196.
- [25] R. SIMION AND F. SCHMIDT, Restricted permutations, *Europ. J. Combin.* **6** (1985) 383–406.

- [26] N. SLOANE AND S. PLOUFFE, The Encyclopedia of Integer Sequences, Academic Press, New York, 1995.
- [27] R. STANLEY, Enumerative Combinatorics, Volume **1**, Cambridge University Press, Cambridge, 1997.
- [28] R. STANLEY, Enumerative Combinatorics, Volume **2**, Cambridge University Press, Cambridge, 1999.
- [29] K. WEGSCHAIDER AND A. RIESE, A Mathematica package for proving hypergeometric multi-sum identities, Available in <http://www.risc.uni-linz.ac.at/research/combinat/risc/software>.
- [30] J. WEST, Permutations with forbidden subsequences and stack sortable permutations, Ph.D. Thesis, M.I.T., (1990).
- [31] H. WILF, The patterns of permutations, *Discr. Math.* **257** (2002) 575–583.
- [32] J. WIMP AND D. ZEILBERGER, Resurrecting the asymptotics of linear recurrences, *J. Math. Anal. Appl.* **111** (1985) 162–176.

MATEMATIK, CHALMERS TEKNISKA HÖGSKOLA OCH GÖTEBORGS UNIVERSITET, S-412 96 GÖTEBORG, SWEDEN.
E-mail address: branden@math.chalmers.se

DEPARTMENT OF MATHEMATICS, HAIFA UNIVERSITY, 31905 HAIFA, ISRAEL.
E-mail address: toufik@math.haifa.ac.il