

An Extension of the Foata Map to Standard Young Tableaux

J. Haglund and L. Stevens

ABSTRACT. We define an inversion statistic on standard Young tableaux. We prove that this statistic has the same distribution over $SYT(\lambda)$ as the major index statistic by exhibiting a bijection on $SYT(\lambda)$ in the spirit of the Foata map on permutations.

RÉSUMÉ. Nous définissons un inversion statistique sur de standard Young tableaux. Nous montrons que cette statistique a la meme distribution au-dessus de $SYT(\lambda)$ que la major index statistique en exhibant un bijection sur $SYT(\lambda)$ dans l'esprit de Foata map sur des permutations.

1. Introduction

A permutation statistic is a combinatorial rule which associates an element of \mathbb{N} to each element of the symmetric group S_n . Let $\sigma = \sigma_1\sigma_2\cdots\sigma_n$ denote the element of S_n which sends i to σ_i for $1 \leq i \leq n$. An inversion of σ is a pair (i, j) , $1 \leq i < j \leq n$ such that $\sigma_i > \sigma_j$. A descent of σ is an integer i , $1 \leq i \leq n - 1$, for which $\sigma_i > \sigma_{i+1}$. The inversion statistic $\text{inv}(\sigma)$ is defined to be the number of inversions of σ and the major index statistic $\text{maj}(\sigma)$ is defined to be the sum of the descents of σ , i.e.

$$\text{inv}(\sigma) = \sum_{i < j, \sigma_i > \sigma_j} 1, \quad \text{maj}(\sigma) = \sum_{i, \sigma_i > \sigma_{i+1}} i.$$

Major P. MacMahon [**MacM**] introduced the major index statistic and proved that, remarkably, its distribution over S_n is equal to the distribution of the inversion statistic over S_n . This raised the question of constructing a canonical bijection $\phi : S_n \rightarrow S_n$ such that $\text{maj}(\sigma) = \text{inv}(\phi(\sigma))$. Foata [**Foa68**] found such a map. (See also [**FoSc78**]). Note that the previous two statistics can be defined on each class \tilde{C} of permutations of a sequence (with repetitions) $1^{m_1}2^{m_2}\cdots r^{m_r}$. The equidistribution of inv and maj also holds for such a class \tilde{C} , a result proved by MacMahon [**MacM**] and reproved by means of the Foata map [**Foa68**], when adequately defined on \tilde{C} . In this paper, we only need the case of permutations without repetitions.

There is a version of maj for tableau which plays a prominent role in symmetric function theory (see for example [**Sta**, Chapter 7] or [**Mac**, Chapter 1]). One could also ask whether there is a natural version of inv which would play a similar role. In this article we advance a candidate tableau inversion statistic, which is defined in terms of a generalization of Foata's map. The statistic was discovered in the course of studying Macdonald polynomials (see the remark at the end of Section 4).

In section one, we review the algorithm describing the Foata map and the necessary background on tableaux. In section two we define the inversion statistic Inv and maps on tableaux that will feature in our Foata-type map. In section three we introduce our Foata-type map and prove that it is a bijection which sends a standard Young tableau with a given Inv to a standard Young tableau with the same major index. Section 5 contains some remarks about extending our map to skew shapes, and in what sense it generalizes Foata's map.

2000 *Mathematics Subject Classification.* 05E10.

Key words and phrases. permutation statistic, tableau statistic, symmetric functions.

J. Haglund supported in part by NSA grant MSPF-02G-193 and NSF grant DMS-0553619.

2. Definitions and Background

We begin by reviewing Foata's map $\phi : S_n \rightarrow S_n$. It can be described as follows. If $n \leq 2$, $\phi(\sigma) = \sigma$. If $n > 2$, we add numbers to ϕ one at a time: begin by setting $\phi^{(1)} = \sigma_1$, $\phi^{(2)} = \sigma_1\sigma_2$. To find $\phi^{(3)}$, start with $\sigma_1\sigma_2\sigma_3$. Then if $\sigma_3 > \sigma_2$, draw a bar after each element of $\sigma_1\sigma_2\sigma_3$ which is less than σ_3 , while if $\sigma_3 < \sigma_2$, draw a bar after each element of $\sigma_1\sigma_2\sigma_3$ which is greater than σ_3 . Also add a bar before σ_1 . For example, if $\sigma = 4137562$, we now have $|41|3$. Now regard the numbers between two consecutive bars as "blocks", and in each block, move the last element to the beginning, and finally remove all of the bars. We end up with $\phi^{(3)} = 143$.

Proceeding inductively, we begin by adding σ_i to the end of $\phi^{(i-1)}$. Then if $\sigma_i > \sigma_{i-1}$, draw a bar after each element of $\phi^{(i-1)}$ which is less than σ_i , while if $\sigma_i < \sigma_{i-1}$, draw a bar after each element of $\phi^{(i-1)}$ which is greater than σ_i . Also draw a bar before $\phi_1^{(i-1)}$. Then in each block, move the last element to the beginning, and finally remove all of the bars. If $\sigma = 4137562$, the successive stages of the algorithm yield

$$\begin{aligned} 143 &= \phi^{(3)} \\ |1|4|3|7 &\mapsto 1437 = \phi^{(4)} \\ |1437|5 &\mapsto 71435 = \phi^{(5)} \\ |71|4|3|5|6 &\mapsto 174356 = \phi^{(6)} \\ |17|4|3|5|6|2 &\mapsto 7143562 = \phi^{(7)}, \end{aligned}$$

so $\phi(4137562) = 7143562$. Note that $\text{maj}(4137562) = 11 = \text{inv}(7143562)$.

THEOREM 2.1. [Foa68] *The map $\phi : S_n \rightarrow S_n$ is a bijection and for all $\sigma \in S_n$, $\text{maj}(\sigma) = \text{inv}(\phi(\sigma))$.*

The inverse to Foata's map is as follows. Let $\psi = \phi^{-1}(\sigma)$. If $\sigma_n > \sigma_1$, draw a bar before each number in σ which is less than σ_n and also before σ_n . If $\sigma_n < \sigma_1$, draw a bar before each number in σ which is greater than σ_n and also before σ_n . Next move each number at the beginning of a block to the end of the block. Remove the bars to obtain $\psi^{(1)}$. The last letter of ψ is now fixed. Now compare $\psi_{n-1}^{(1)}$ with $\psi_1^{(1)}$. Create blocks as above, drawing a bar before $\psi_{n-1}^{(1)}$. Proceed in this manner. For example, if $\sigma = 7143562$, the successive stages of the algorithm yield

$$\begin{aligned} |71|4|3|5|6|2 &\mapsto 1743562 = \psi^{(1)} \\ |17|4|3|5|6|2 &\mapsto 7143562 = \psi^{(2)} \\ |7143|562 &\mapsto 1437562 = \psi^{(3)} \\ |1|4|3|7562 &\mapsto 1437562 = \psi^{(4)} \\ |14|37562 &\mapsto 4137562 = \psi^{(5)} \\ 4137562 &= \psi^{(6)} = \psi^{(7)}, \end{aligned}$$

so $\phi^{-1}(7143562) = 4137562$. It turns out that the generalization of ϕ^{-1} to tableaux is simpler to describe than the generalization of ϕ .

Next we review the necessary background on tableaux. A partition of a positive integer n is a sequence $\lambda = (\lambda_1, \dots, \lambda_k)$ of positive integers such that $\sum_{i=1}^k \lambda_i = n$ and $\lambda_1 \geq \dots \geq \lambda_k$. If λ is a partition of n we write $\lambda \vdash n$ or $|\lambda| = n$. The λ_i are called the parts of λ .

We represent $\lambda = (\lambda_1, \dots, \lambda_k)$ pictorially as follows. We assign (row, column)-coordinates to unit squares in the first quadrant, obtained by permuting the (x, y) coordinates of the upper right-hand corner of the square, so the lower left-hand square has coordinates $(1, 1)$, the square above it $(2, 1)$, etc., and a square (i, j) has ordinate i and abscissa j . Then the Ferrer's diagram of λ , which we also denote by λ , is the set of squares, or "cells" $\{(i, j) \in \mathbb{N} \times \mathbb{N} : 1 \leq i \leq k, 1 \leq j \leq \lambda_i\}$. That is, the diagram of λ consists of k left-justified rows of squares in the first quadrant of the xy -plane with λ_i squares in the i th row from the bottom. See Figure 1.




FIGURE 1. The Ferrer's diagram of the partition $(4, 3, 1)$

The conjugate partition of $\lambda = (\lambda_1, \dots, \lambda_k)$ is $\lambda' = (\lambda'_1, \dots, \lambda'_j)$, where λ'_i is the number of parts of λ which are greater than or equal to i . Geometrically, the diagram of λ' is the reflection of the diagram of λ about the line $y = x$.

If $\lambda \vdash n$, a standard Young tableau of shape λ is a filling of the diagram of λ with the numbers $1, \dots, n$ such that in each row, the numbers are increasing from left to right, and in each column, the numbers are increasing from bottom to top. The number filling a cell is called the content of the cell. We let $c(i, j)$ denote the content of the cell (i, j) .

We let $SYT(\lambda)$ denote the set of standard Young tableaux of shape λ . For $T \in SYT(\lambda)$, the conjugate tableau $T' \in SYT(\lambda')$ is obtained by filling the cell (i, j) with the content of the cell (j, i) of T .

3. An Inv Statistic on Standard Young Tableaux

For a standard Young tableau T , the major index of T is given by

$$\text{maj}(T) = \sum_{i \in \text{Des}(T)} i,$$

where $\text{Des}(T) = \{i \mid i + 1 \text{ is in a row above the row containing } i \text{ in } T\}$. Here we define another statistic on standard Young tableaux which we call the Inv statistic. Let $(i, j), (k, l) \in T$ such that $c(i, j) > c(k, l)$. The cells (i, j) and (k, l) form an “inversion pair” if (k, l) is weakly SE of (i, j) , and they do not form an inversion pair if (k, l) is weakly NW of (i, j) . If (k, l) is strictly SW of (i, j) , then whether or not they form an inversion pair is determined by a path in T called the “inversion path” of (i, j) . By strictly SW, we mean Southwest but not due South or due West, i.e. $k < i$ and $l < j$. Below, we construct the set of inversion paths for T . In the construction we define maps on standard Young tableaux which will be used in our extension of the Foata map.

DEFINITION 3.1. Let $T \in SYT(\lambda)$, where $\lambda \vdash n$. For $k = 1, \dots, n$, let $\pi(T, k)$ be the path in T constructed according to the following algorithm:

Start at the lower left-hand corner of the cell with content k . Suppose this cell has coordinates (i, j) . If $i = 1$ or $j = 1$, proceed in a straight line until you reach the origin. If not, compare the contents of the cells $(i - 1, j)$ and $(i, j - 1)$. If $c(i - 1, j) > c(i, j - 1)$, take one unit step South. If $c(i - 1, j) < c(i, j - 1)$, take one unit step West. Arriving at the lower left-hand corner of a new cell (i', j') , iterate the algorithm. Proceed until you reach the origin.

For example, let T be the SYT in Figure 2 (the fact that the shape is a square is coincidental). To form the inversion path $\pi(T, 16)$, we start at the lower left-hand corner of the square containing 16, and draw a unit line segment left, since the square to the left of 16 contains 15, which is greater than the number in the square below (namely 14). Next we compare the numbers in the squares to the left and below the square containing 15. The largest is 13, so we now extend our inversion path by drawing another unit line segment, in this case downwards, ending at the lower left-hand corner of the square containing 13. We continue comparing squares left and below, moving one unit in the direction of the largest, until we reach the edge of the tableau (the lower left-hand corner of the square containing 3), at which time we go in the only direction we can, namely straight down, until we reach a square with no left or bottom neighbors (the square containing 1).

Note that the path $\pi(T, k)$ partitions the cells of T with content less than k into two sets: cells which are weakly NW of $\pi(T, k)$ and cells which are weakly SE of $\pi(T, k)$. We will say that cells weakly NW of $\pi(T, k)$ are above $\pi(T, k)$ and cells weakly SE of $\pi(T, k)$ are below $\pi(T, k)$. Two cells are on the same side of $\pi(T, k)$ if they are both above or both below $\pi(T, k)$. Note also that since $\pi(T, k)$ is

6	10	15	16
5	9	13	14
3	7	11	12
1	2	4	8

FIGURE 2. A standard Young tableau T and the path $\pi(T, 16)$

6	10	15	16
5	9	13	14
3	7	11	12
1	2	4	8

 \longrightarrow

5	9	14	16
4	8	13	15
2	6	11	12
1	3	7	10

FIGURE 3. The tableau on the right is $\Psi_{16}(T)$

constructed taking only South and West steps from the cell with content k , all cells weakly SE of the cell with content k are always below $\pi(T, k)$ and all cells weakly NW of the cell with content k are always above $\pi(T, k)$.

DEFINITION 3.2. Let $T \in SYT(\lambda)$, where $\lambda \vdash n$. For $k = 1, \dots, n$, let $\Psi_k(T)$ be the tableau constructed according to the following algorithm:

Construct the path $\pi(T, k)$. Partition the cells of T with content less than k into “blocks” $\{(i_1, j_1), \dots, (i_m, j_m)\}$ of maximal length such that

- (i) $c(i_l, j_l) = c(i_{l+1}, j_{l+1}) - 1$ for $1 \leq l \leq m - 1$.
- (ii) (i_1, j_1) is on the same side of $\pi(T, k)$ as $(1, 1)$, and (i_l, j_l) is on the other side of $\pi(T, k)$ for $2 \leq l \leq m$,

Note that blocks may consist of only one cell. Next perform the following “cycling” procedure on the contents of the cells in each block:

- (i) Replace $c(i_1, j_1)$ by $c(i_m, j_m)$.
- (ii) For $2 \leq l \leq m$, replace $c(i_l, j_l)$ by $c(i_l, j_l) - 1$.

The resulting tableau is $\Psi_k(T)$.

In the next section, we prove that $\Psi_k(T) \in SYT(\lambda)$. Note that the maps Ψ_2, Ψ_1 are the identity.

Simply put, the blocks for the path $\pi(T, k)$ consist of cells containing consecutive numbers such that the smallest number is in a cell on the same side of the path as the cell containing 1 and all of the rest of the cells are on the other side of the path. In the example in Figure 2, if we denote cells by their contents, then the blocks for $\pi(T, 16)$ are $\{1\}$, $\{2, 3\}$, $\{4, 5, 6, 7\}$, $\{8, 9, 10\}$, $\{11\}$, $\{12\}$, $\{13\}$, $\{14, 15\}$.

If the contents of the cells in a block are $a, a + 1, \dots, a + j$, then the cycling procedure sends a to the cell which had content $a + 1$, etc. and finally $a + j$ to the cell which had content a . See Figure 3. A key observation about the cycling procedure is that it preserves the relative order of the contents of cells in different blocks.

DEFINITION 3.3. The collection of paths

$$\{\pi(T, n), \pi(\Psi_n(T), n - 1), \dots, \pi(\Psi_3 \circ \dots \circ \Psi_n(T), 2)\}$$

is the set of inversion paths for T . For $(i, j) \in T$, $(i, j) \neq (1, 1)$, the inversion path of (i, j) is the unique element of the set of inversion paths for T starting at the lower left-hand corner of the cell (i, j) .

In our running example, the path $\pi(\Psi_{16}(T), 15)$ is the inversion path for the cell $(3, 4)$ (the cell with content 14) in T . See Figure 4. See also Figures 6 and 7 for a complete example.

Now we are ready to introduce the Inv statistic.

5	9	14	16
4	8	13	15
2	6	11	12
1	3	7	10

6	10	15	16
5	9	13	14
3	7	11	12
1	2	4	8

FIGURE 4. On the left: $\Psi_{16}(T)$ and the path $\pi(\Psi_{16}(T), 15)$. On the right: T and the inversion path of the cell $(3, 4)$.

DEFINITION 3.4. Let $T \in SYT(\lambda)$. An ordered pair of cells $((i, j), (k, l))$ in T is an inversion pair if $c(k, l) < c(i, j)$ and (k, l) is below the inversion path of (i, j) . $\text{Inv}(T)$ is the total number of inversion pairs in T .

In our running example, the cell with content 16 in T forms inversion pairs with the cells with contents 1, 2, 4, 8, 11, 12, 13, 14 and the cell with content 14 in T forms inversion pairs with the cells with contents 4, 8, 11, 12.

4. The bijection

Let $\lambda \vdash n$, and let $\Psi = \Psi_3 \circ \dots \circ \Psi_n$. One of our main results is:

THEOREM 4.1. *The map $\Psi : SYT(\lambda) \rightarrow SYT(\lambda)$ is a bijection and for $T \in SYT(\lambda)$, $\text{Inv}(T) = \text{maj}(\Psi(T))$.*

Most of this section is devoted to outlining the proof of Theorem 4.1. First one proves that $\Psi_k : SYT(\lambda) \rightarrow SYT(\lambda)$. One uses the following lemma, which helps us understand the geometry of the blocks for a given path.

LEMMA 4.2. *Let $T \in SYT(\lambda)$. Suppose that the cells with contents $a, a+1, \dots, a+m$ form a block for $\pi(T, k)$. If the cell $(1, 1)$ is below $\pi(T, k)$, then the cells containing $a+1, \dots, a+m$ are strictly NW of the cell containing a . If the cell $(1, 1)$ is above $\pi(T, k)$, then the cells containing $a+1, \dots, a+m$ are strictly SE of the cell containing a .*

THEOREM 4.3. *For $k = 3, \dots, n$, if $T \in SYT(\lambda)$, then $\Psi_k(T) \in SYT(\lambda)$.*

Next one shows that the maps Ψ_k are bijections by explicitly constructing the inverse maps. Our starting point is the following lemma, which is key in reconstructing the path $\pi(\Psi_k^{-1}(S), k)$.

LEMMA 4.4. *Let $T \in SYT(\lambda)$, where $\lambda \vdash n$. For $k = 2, \dots, n$, $(1, 1)$ is below $\pi(T, k)$ if and only if $k-1 \in \text{Des}(\Psi_k(T))$.*

Now we are ready to describe the maps $\Phi_k = \Psi_k^{-1}$. Let $S \in SYT(\lambda)$, and suppose $c(r, s) = k$ and $c(x, y) = k-1$. We give an algorithm to obtain a new tableau from S and a path $\tilde{\pi}(S, k)$ starting at the lower left-hand corner of the cell (r, s) and ending at the origin:

Step 1: If $r = 1$ or $s = 1$, draw a straight line to the origin and stop. Otherwise:

If $k-1 \in \text{Des}(S)$: Find all “simple blocks” $\{(i_1, j_1), \dots, (i_m, j_m)\}$ of cells such that

- (i) $c(r-1, s-1) < c(i_l, j_l) < k$ and $c(i_l, j_l) = c(i_{l+1}, j_{l+1}) + 1$ for all $1 \leq l \leq m$,
- (ii) (i_1, j_1) is weakly SE of (r, s) and $(i_2, j_2), \dots, (i_m, j_m)$ are weakly NW of (r, s) ,
- (iii) the cell with content $c(i_m, j_m) - 1$ is weakly SE of (r, s) .

If $k-1 \notin \text{Des}(S)$: Find all simple blocks $\{(i_1, j_1), \dots, (i_m, j_m)\}$ of cells such that

- (i) $c(r-1, s-1) < c(i_l, j_l) < k$ and $c(i_l, j_l) = c(i_{l+1}, j_{l+1}) + 1$ for all $1 \leq l \leq m$,
- (ii) (i_1, j_1) is weakly NW of (r, s) and $(i_2, j_2), \dots, (i_m, j_m)$ are weakly SE of (r, s) ,
- (iii) the cell with content $c(i_m, j_m) - 1$ is weakly NW of (r, s) .

Step 2: Perform the following “reverse cycling” procedure on the contents of the cells of each block from Step 1: Replace $c(i_1, j_1)$ by $c(i_m, j_m)$, and for $2 \leq l \leq m$, replace $c(i_l, j_l)$ by $c(i_l, j_l) + 1$.

Comment: In Step 1, we are looking for the blocks of $\pi(\Psi_k^{-1}(S), k)$ which are easy to identify, in particular those which are independent of the path. In Step 2, we undo the cycling procedure of Ψ_k on these blocks. To identify the rest of the blocks, we must reconstruct the path.

Step 3: Construct a unit segment of $\tilde{\pi}(S, k)$ as follows:

If $k - 1 \in \text{Des}(S)$: Go South if $(r - 1, s)$ is in one of the blocks from Step 2 and $c(r - 1, s) > c(r, s - 1)$. Otherwise go West.

If $k - 1 \notin \text{Des}(S)$: Go West if $(r, s - 1)$ is in one of the blocks from Step 2 and $c(r, s - 1) > c(r - 1, s)$. Otherwise go South.

Step 4: A portion of $\tilde{\pi}(S, n)$ has been constructed and ends at the lower left-hand corner of the cell (u, v) . If $u = 1$ or $v = 1$, draw a straight line to the origin and stop. Otherwise:

Find all simple blocks $\{(i_1, j_1), \dots, (i_m, j_m)\}$ of cells such that

- (i) for all $1 \leq k \leq m$, (i_k, j_k) was not in a simple block at any previous stage in the algorithm,
- (ii) $c(u - 1, v - 1) < c(i_l, j_l) < k$ and $c(i_l, j_l) = c(i_{l+1}, j_{l+1}) + 1$ for all $1 \leq k \leq m$,
- (iii) (i_1, j_1) and (x, y) are on the same side of the portion of the path constructed so far and $(i_2, j_2), \dots, (i_m, j_m)$ are on the other side,
- (iv) the cell with content $c(i_m, j_m) - 1$ and (i_1, j_1) are on the same side of portion of the path constructed so far.

Step 5: Reverse cycle the contents of the cells of each block from Step 4.

Step 6: Construct a unit segment of $\tilde{\pi}(S, k)$ as follows:

If $n - 1 \in \text{Des}(S)$: Go South if $(u - 1, v)$ has already appeared in a simple block and $c(u - 1, v) > c(u, v - 1)$. Otherwise go West.

If $n - 1 \notin \text{Des}(S)$: Go West if $(u, v - 1)$ has already appeared in a simple block and $c(u, v - 1) > c(u - 1, v)$. Otherwise go South.

Arriving at the lower left-hand corner of a new cell, iterate the algorithm from Step 4.

Observe that if a cell with content a appears in a simple block, then all cells with content greater than a and less than n must have already appeared in simple blocks. Moreover, upon completion of the algorithm, every cell with content greater than 1 and less than n will appear in exactly one simple block.

In Figure 5, we give an example of the algorithm with $k = 16$ applied to the tableau $\Psi_{16}(T)$ from Figure 3.

THEOREM 4.5. *For $k = 3, \dots, n$, if T is the tableau obtained by applying the above algorithm to the tableau S , then $S = \Psi_k(T)$.*

In Figure 6, we give an example of a tableau S and its image under the map Ψ . We have $\text{maj}(\Psi(S)) = 1 + 4 + 5 + 7 = 17$. In Figure 7, we show the set of inversion paths for S . If we label cells by their contents, then inversion pairs for S are $(8, 6), (8, 5), (8, 4), (8, 3), (8, 2), (8, 1), (7, 6), (7, 5), (7, 4), (7, 3), (7, 2), (7, 1), (6, 5), (4, 1), (3, 2), (3, 1)$. So $\text{Inv}(S) = 17$.

COROLLARY 4.6. *The statistics maj and Inv have equal distributions over $\text{SYT}(\lambda)$.*

Remark: Part of our motivation for defining an inversion statistic on SYT is based on recent work of Haglund, Haiman and Loehr [HHL], where it is shown that the coefficient of a monomial symmetric function in the Macdonald polynomial $\tilde{H}_\mu[X; q, t]$ can be expressed in terms of descents and inversion pairs for fillings of the Ferrers shape of μ by positive integers. One could hope that a similar phenomenon would apply to the coefficients $\tilde{K}_{\lambda, \mu}(q, t)$ in the expansion of \tilde{H}_μ into Schur functions (Macdonald showed that $\tilde{K}_{\lambda, \mu}(1, 1)$ equals the number of $\text{SYT}(\lambda)$, and posed the question [Mac, p. 356] of whether $\tilde{K}_{\lambda, \mu}(q, t)$ could be written in the form

$$(4.1) \quad \tilde{K}_{\lambda, \mu}(q, t) = \sum_{T \in \text{SYT}(\lambda)} t^{\text{tstat}(T, \mu)} q^{\text{qstat}(T, \mu)}$$

for some tableau statistics $\text{tstat}, \text{qstat}$). In [Hag04, Conjecture 3] specific values for $\text{tstat}, \text{qstat}$ are conjectured for all μ with at most three columns. The value of tstat is described in terms of descents of T , while the value of qstat is described in terms of certain ‘‘inversion pairs’’ of T . (The special case of this conjecture where μ has at most two columns is proved in [HHL, Proposition 9.2]). Although we




FIGURE 5. An example of the map Φ_n




FIGURE 6. On the left: a tableau S . On the right: $\Psi(S)$.




FIGURE 7. The set of inversion paths for S .

have as yet been unable to extend this conjecture to general μ , Corollary 4.6 gives a way of expressing $\tilde{K}_{\lambda,(n)}(q,t)$ in this form, since it is known that

$$(4.2) \quad \tilde{K}_{\lambda,(n)}(q,t) = \sum_{T \in SYT(\lambda)} q^{\text{maj}(T)},$$

which by Corollary 4.6 also equals

$$(4.3) \quad \tilde{K}_{\lambda,(n)}(q,t) = \sum_{T \in SYT(\lambda)} q^{\text{Inv}(T)}.$$

5. Extensions

Although we have assumed throughout this article that λ is a partition shape, all our results apply just as easily to skew shapes. For a given SYT T of skew shape $\lambda \vdash n$, note that our map Φ (and its inverse) both fix the largest element n in the tableau. Thus our inversion statistic is equidistributed with maj over the set of $SYT(\lambda)$ with n occurring in some fixed corner square. Let

$$\text{comaj}(T) = \sum_{i \in \text{Des}(T)} n - i.$$

It is known that comaj and maj have the same distribution over SYT of skew shape [Sta, Chapter 7]. We briefly describe a parallel version of our Foata map which is linked to comaj . If square (i,j) contains 1 in T , start at the NE corner of (i,j) and draw a unit line segment N if $c(i+1,j) > c(i,j+1)$, otherwise draw a unit line segment E , and now iterate, eventually ending up at the upper border of T . Next break up the numbers 2 through n into maximal blocks, as before, consisting of an integer k on the same side of the inversion path as n , and a consecutive sequence of integers $k-1, k-2, \dots, k-a$ on the other side of the path, then cycling by moving $k-a$ to the square containing k , k to the square containing $k-1$, etc.. Thus each square ends up with an inversion path which starts at its NE corner and travels NE to the border, and the number of inversion pairs are equidistributed with comaj over the set of all $SYT(\lambda)$ with 1 occurring in some fixed corner square.

We now show that the special case of our map Φ when λ is a disjoint union of squares, i.e. $\lambda = (n, n-1, \dots, 1)/(n-1, \dots, 1)$, and T is the tableau obtained by filling these squares with the numbers $\sigma_1, \dots, \sigma_n$ (see Figure 8), is essentially the same as Foata's original map ϕ applied to σ^{-1} . We identify T with σ and use the notation $\Phi(\sigma)$. To find $\Phi_k(\sigma)$, partition the numbers $k-1, \dots, 1$ into blocks $\{a, a-1, \dots, a-j\}$ of maximal length such that a is on the same side of k as $k-1$ and $a-1, \dots, a-j$ are on the other side of k . Then reverse cycle, e.g. $\Phi_4(346251) = 146352$. Let $\sigma^{-1} = \omega$, so ω_i is the position of i in σ . We now show that $\phi(\omega) = (\Phi(\sigma))^{-1}$. The main idea is to prove that for any k , $3 \leq k \leq n$, $\phi^{(k)}\omega_{k+1} \dots \omega_n = (\Phi_k \circ \dots \circ \Phi_3(\sigma))^{-1}$, where $\phi^{(k)} = \phi^{(k)}(\omega)$. This is straightforward when $k=3$; $\Phi_3(\sigma) = \sigma$ unless 2 and 1 are on opposite sides of 3, in which case Φ_3 interchanges their positions and leaves all other numbers fixed, whereas $\phi^{(3)} = \omega_1\omega_2\omega_3$ only if ω_1 and ω_2 are both less than or greater than ω_3 , otherwise $\phi^{(3)} = \omega_2\omega_1\omega_3$. Now assume that the claim is true for some k such that $3 \leq k \leq n-1$. Write $\Phi_k \circ \dots \circ \Phi_3(\sigma) = \beta$ and $\phi^{(k)} = \alpha_1 \dots \alpha_k$. Then $\{a, a-1, \dots, a-j\}$ is a block in β if and only if $\alpha_{a-j} \dots \alpha_{a-1}\alpha_a$ is a block in $\alpha_1 \dots \alpha_k\omega_{k+1}$. The reverse cycling procedure sends $a-j$ to the position of a and $a, \dots, a-j+1$ to the positions of $a-1, \dots, a-j$, respectively, which corresponds to the Foata algorithm sending the block $\alpha_{a-j} \dots \alpha_{a-1}\alpha_a$ to $\alpha_a\alpha_{a-j} \dots \alpha_{a-1}$. Thus $\phi^{(k+1)}\omega_{k+1} \dots \omega_n = (\Phi_{k+1}(\beta))^{-1}$. For the example in Figure 8, the sequence $\sigma, \Phi_3(\sigma), \Phi_4 \circ \Phi_3(\sigma), \Phi_5 \circ \Phi_4 \circ \Phi_3(\sigma), \Phi(\sigma)$ is

$$346251, 346251, 146352, 146253, 256143$$

whereas the sequence $\omega, \phi^{(3)}\omega_4\omega_5\omega_6, \phi^{(4)}\omega_5\omega_6, \phi^{(5)}\omega_6, \phi(\omega)$ is

$$641253, 641253, 164253, 146253, 416523.$$

6. Concluding Remarks

The authors would like to thank the referees for helpful suggestions on the exposition, as well as pointing out some interesting directions for future research. For example, MacMahon's result on the equidistribution of maj and inv holds for multiset permutations, and Foata showed how his map proves this more general fact bijectively. Perhaps our results on SYT also have versions for SSYT. We remark




FIGURE 8. On the left: a tableau T of shape $(6, 5, 4, 3, 2, 1)/(5, 4, 3, 2, 1)$ whose filling corresponds to the permutation $\sigma = 346251$. On the right: $\Phi(T)$.

that the obvious thing to try, which is to start with a SSYT, standardize in the usual way, apply the Φ map, then “unstandardize”, doesn’t quite work, as this process will in general produce fillings which are not column strict, and hence not SSYT. One thing that does seem to have a natural analogue in our setting is that of the “inversion code” of a permutation, which is a sequence $x_1 x_2 \dots x_n$ with $0 \leq x_i \leq i - 1$ for $1 \leq i \leq n$, where x_i equals the number of inversion pairs of the permutation of the form (i, j) with $i > j$. Our tableaux inversion statistic is exactly the number of similar such pairs, as in the example just above Corollary 4.6. One could also hope to connect our ideas with other inversion statistics which occur in the theory of symmetric functions and Macdonald polynomials, such as the inversion statistic for tuples of SSYT which can be used to describe LLT polynomials (see [HHL]).

References

- [Foa68] D. Foata, *On the Netto inversion number of a sequence*, Proc. Amer. Math. Soc. **19** (1968), 236–240.
- [FoSc78] D. Foata and M. P. Schützenberger, *Major index and inversion number of permutations*, Math. Nachr., **83** (1978), 143–159.
- [Hag04] J. Haglund, *A combinatorial model for the Macdonald polynomials*, Proc. Nat. Acad. Sci. U.S.A., **101** (2004), 16127–16131.
- [HHL] J. Haglund, M. Haiman, and N. Loehr, *A combinatorial formula for Macdonald polynomials*, Jour. Amer. Math. Soc., **18** (2005), 735–761.
- [Mac] I. G. Macdonald, *Symmetric Functions and Hall Polynomials*, Oxford Mathematical Monographs, second ed., Oxford Science Publications, The Clarendon Press Oxford University Press, New York, 1995.
- [MacM] Major P. A. MacMahon, *Two applications of general theorems in combinatory analysis*, Proc. London math. Soc., **15** (1916), 314–321.
- [Sta] R. P. Stanley, *Enumerative Combinatorics*, Cambridge University Press, Cambridge, United Kingdom, Volume 2, 1999.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF PENNSYLVANIA, PHILADELPHIA, PA 19104-6395

E-mail address: `jhaglund@math.upenn.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, SAN DIEGO, LA JOLLA, CA 92093-0112

E-mail address: `stevens@math.ucsd.edu`