

On the diagonal ideal of $(\mathbb{C}^2)^n$ and q, t -Catalan numbers

Kyungyong Lee^{1†} and Li Li²

¹Department of Mathematics, Purdue University, West Lafayette, IN 47907

²Department of Mathematics, University of Illinois at Urbana Champaign, Urbana, IL 61801

Abstract. Let I_n be the (big) diagonal ideal of $(\mathbb{C}^2)^n$. Haiman proved that the q, t -Catalan number is the Hilbert series of the graded vector space $M_n = \bigoplus_{d_1, d_2} (M_n)_{d_1, d_2}$ spanned by a minimal set of generators for I_n . We give simple upper bounds on $\dim (M_n)_{d_1, d_2}$ in terms of partition numbers, and find all bi-degrees (d_1, d_2) such that $\dim (M_n)_{d_1, d_2}$ achieve the upper bounds. For such bi-degrees, we also find explicit bases for $(M_n)_{d_1, d_2}$.

Résumé. Soit I_n l'idéal de la (grande) diagonale de $(\mathbb{C}^2)^n$. Haiman a démontré que le q, t -nombre de Catalan est la série de Hilbert de l'espace vectoriel gradué $M_n = \bigoplus_{d_1, d_2} (M_n)_{d_1, d_2}$ engendré par un ensemble minimal de générateurs de I_n . Nous obtenons des bornes supérieures simples pour $\dim (M_n)_{d_1, d_2}$ en termes de nombres de partitions, ainsi que tous les bi-degrés (d_1, d_2) pour lesquels ces bornes supérieures sont atteintes. Pour ces bi-degrés, nous trouvons aussi des bases explicites de $(M_n)_{d_1, d_2}$.

Keywords: q, t -Catalan number, diagonal ideal

1 introduction

1.1 Background

The goal of this paper is to study the q, t -Catalan numbers and the (thick) diagonal ideal in $(\mathbb{C}^2)^n$, and discuss some technique that we have developed recently.

Let n be a positive integer. Consider the set of n -tuples $\{(x_i, y_i)\}_{1 \leq i \leq n}$ in the plane \mathbb{C}^2 . They form an affine space $(\mathbb{C}^2)^n$ with coordinate ring $\mathbb{C}[\mathbf{x}, \mathbf{y}] = \mathbb{C}[x_1, y_1, \dots, x_n, y_n]$. There is a natural symmetric group S_n acting on $\mathbb{C}[\mathbf{x}, \mathbf{y}]$ by permuting the coordinates in \mathbf{x}, \mathbf{y} simultaneously. With this group action, a polynomial $f \in \mathbb{C}[\mathbf{x}, \mathbf{y}]$ is called *alternating* if

$$\sigma(f) = \text{sgn}(\sigma)f \quad \text{for all } \sigma \in S_n.$$

Define $\mathbb{C}[\mathbf{x}, \mathbf{y}]^\epsilon$ to be the vector space of alternating polynomials in $\mathbb{C}[\mathbf{x}, \mathbf{y}]$.

[†]Research of the first author partially supported by NSF grant DMS 0901367.

There is a more combinatorial way to describe $\mathbb{C}[\mathbf{x}, \mathbf{y}]^\epsilon$. Denote by \mathbb{N} the set of nonnegative integers. Let \mathfrak{D}_n be the set of n -tuples $D = \{(\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n)\} \subset \mathbb{N} \times \mathbb{N}$. For $D \in \mathfrak{D}_n$, define

$$\Delta(D) := \det \begin{bmatrix} x_1^{\alpha_1} y_1^{\beta_1} & x_1^{\alpha_2} y_1^{\beta_2} & \dots & x_1^{\alpha_n} y_1^{\beta_n} \\ \vdots & \vdots & \ddots & \vdots \\ x_n^{\alpha_1} y_n^{\beta_1} & x_n^{\alpha_2} y_n^{\beta_2} & \dots & x_n^{\alpha_n} y_n^{\beta_n} \end{bmatrix}$$

Then $\{\Delta(D)\}_{D \in \mathfrak{D}_n}$ forms a basis for the \mathbb{C} -vector space $\mathbb{C}[\mathbf{x}, \mathbf{y}]^\epsilon$.

It is easy to see that any alternating polynomial vanishes on the thick diagonal of $(\mathbb{C}^2)^n$. (By thick diagonal we mean the set of n -tuples of points in \mathbb{C}^2 where at least two points coincide.) A theorem of Haiman asserts that the converse is also true: any polynomial that vanishes on the diagonal of $(\mathbb{C}^2)^n$ can be generated by alternating polynomials, i.e.

$$\bigcap_{1 \leq i < j \leq n} (x_i - x_j, y_i - y_j) = \text{ideal generated by } \Delta(D)\text{'s.}$$

We call the above ideal the **diagonal ideal** and denote it by I_n . the number of minimal generators of I_n , which is the same as the dimension of the vector space $M_n = I_n/(\mathbf{x}, \mathbf{y})I_n$, is equal to the n -th Catalan number. The space M_n is doubly graded as $\bigoplus_{d_1, d_2} (M_n)_{d_1, d_2}$. The q, t -Catalan number, originally introduced by A.M.Garsia and M.Haiman in [4], can be defined as

$$C_n(q, t) = \sum_{d_1, d_2} t^{d_1} q^{d_2} \dim(M_n)_{d_1, d_2}.$$

The q, t -Catalan number $C_n(q, t)$ also has a combinatorial interpretation using Dyck paths. To be more precise, take the $n \times n$ square whose southwest corner is $(0, 0)$ and northeast corner is (n, n) . Let \mathcal{D}_n be the collection of Dyck paths, i.e. lattice paths from $(0, 0)$ to (n, n) that proceed by NORTH or EAST steps and never go below the diagonal. For any Dyck path Π , let $a_i(\Pi)$ be the number of squares in the i -th row that lie in the region bounded by Π and the diagonal. A.M.Garsia and J.Haglund ([2], [3]) among others showed that

$$C_n(q, t) = \sum_{\Pi \in \mathcal{D}_n} q^{\text{area}(\Pi)} t^{\text{dinv}(\Pi)},$$

where

$$\text{dinv}(\Pi) := |\{(i, j) \mid i < j \text{ and } a_i(\Pi) = a_j(\Pi)\}| + |\{(i, j) \mid i < j \text{ and } a_i(\Pi) + 1 = a_j(\Pi)\}|.$$

Haiman posed a question asking for a rule that associate to each Dyck path Π an element $D(\Pi) \in \mathfrak{D}_n$ such that $\deg_x \Delta(D(\Pi)) = \text{area}(\Pi)$, $\deg_y \Delta(D(\Pi)) = \text{dinv}(\Pi)$, and that the set $\{\Delta(D(\Pi))\}$ generates I_n . The last condition is equivalent to requiring the images $\{\overline{\Delta(D(\Pi))}\}$ form a basis of M_n . It is natural to ask the following more general question:

Question 1.1 Given a bi-degree (d_1, d_2) , is there a combinatorially significant construction of the basis for each $(M_n)_{d_1, d_2}$?

1.2 Main results

This paper initiates the approach to the study of M_n by comparing it with $M_{n'}$ for a large integer n' . On the one hand, there is a natural map $M_n \rightarrow M_{n'}$ for any $n' > n$. On the other hand, for n' sufficiently large, the basis of $(M_{n'})_{d'_1, d_2}$ becomes “stable” if we fix d_2 and fix

$$k = \binom{n}{2} - d_1 - d_2 = \binom{n'}{2} - d'_1 - d_2.$$

Therefore we can take the “limit” of such basis for $n' \rightarrow \infty$. This basis is indexed by the partitions of k . As a consequence, $(M_{n'})_{d'_1, d_2}$ can be imbedded as a subspace of the polynomial ring with infinite many variables $\mathbb{C}[\rho_1, \rho_2, \dots]$. The induced map

$$\bar{\varphi} : (M_n)_{d_1, d_2} \rightarrow \mathbb{C}[\rho_1, \rho_2, \dots],$$

which will be defined explicitly in subsection 1.2.3, provides a powerful tool to study M_n .

1.2.1 Asymptotic behavior when $k \ll n$

We shall show that if $k \ll n$, then $(M_n)_{d_1, d_2}$ has a basis $\{\overline{\Delta(D)}\}$ where D are so-called **minimal staircase forms** that will be defined later.

The essential step is to observe the following three linear relations that turn the questions into combinatorial games. First we introduce some notations.


- For $D = \{P_1, \dots, P_n\} \in \mathfrak{D}_n$ where $P_i = (\alpha_i, \beta_i)$, define $|P_i| = \alpha_i + \beta_i$.

Relation 1. Given positive integers $1 \leq i \neq j \leq n$ such that $|P_i| = i - 1, |P_{i+1}| = i, |P_j| = j - 1, |P_{j+1}| = j, \beta_i > 0, \alpha_j > 0$ (we assume $|P_{n+1}| = n$). Let D' be obtained from D by moving P_i to southeast and P_j to northwest, i.e.

$$D' = \{P_1, \dots, P_{i-1}, P_i + (1, -1), P_{i+1}, \dots, P_{j-1}, P_j + (-1, 1), P_{j+1}, \dots, P_n\}.$$

Then $\overline{\Delta(D)} = \overline{\Delta(D')}$.

Example: $n = 9, i = 2, j = 6$.




Relation 2. Given positive integers h, ℓ and m such that $2 \leq h < h + \ell + m \leq n + 1, |P_h| = h - 1, |P_{h+\ell}| = h + \ell - 1, |P_{h+\ell+m}| = h + \ell + m - 1$ (by convention, the last equality holds if $h + \ell + m = n + 1$) and $\alpha_{h+\ell}, \dots, \alpha_{h+\ell+m-1} \geq \ell$. Let D' be obtained from D by moving the m points $P_{h+\ell}, \dots, P_{h+\ell+m-1}$ to the left by ℓ units and moving the ℓ points $P_h, \dots, P_{h+\ell-1}$ to the right by m units, i.e.

$$D' = \{P_1, P_2, \dots, P_{h-1}, P_{h+\ell} - (\ell, 0), P_{h+\ell+1} - (\ell, 0), \dots, P_{h+\ell+m-1} - (\ell, 0), P_h + (m, 0), P_{h+1} + (m, 0), \dots, P_{h+\ell-1} + (m, 0), P_{h+\ell+m}, \dots, P_n\}.$$

Then $\overline{\Delta(D)} = \overline{\Delta(D')}$.

Example: $n = 10, h = 3, \ell = 4, m = 3$.




Relation 3. Given positive integers j and s . Suppose P_{s_0} is the last point in D satisfying $|P_i| = i - 1$. Define $j = (s_0 - 1 - |P_{s_0}|) + (s_0 - |P_{s_0+1}|) + \dots + (n - 1 - |P_n|)$. Suppose $|P_i| = i - 1$ for $1 \leq i \leq j + 2$, $P_2 = (1, 0)$, $s_0 \leq s \leq n$, and $\alpha_s, \beta_s \geq 1$. Let

$$D^{\nwarrow} = \{P_1, \dots, P_{j+1}, P_{j+2} + (1, -1), P_{j+3}, \dots, P_{s-1}, P_s + (-1, 1), P_{s+1}, \dots, P_n\},$$

$$D^{\searrow} = \{P_1, (0, 1), P_3, \dots, P_{s-1}, P_s + (1, -1), P_{s+2}, \dots, P_n\}.$$

Then $2\overline{\Delta(D)} = \overline{\Delta(D^{\nwarrow})} + \overline{\Delta(D^{\searrow})}$.


Example: $n = 9, i = 2, j = 6$.



We call $D = \{P_1, \dots, P_n\}$ a *minimal staircase form* if $|P_i| = i - 1$ or $i - 2$ for every $1 \leq i \leq n$. For a minimal staircase form D , let $\{i_1 < i_2 < \dots < i_\ell\}$ be the set of i 's such that $|P_i| = i - 1$, we define the *partition type* of D to be the partition of $\binom{n}{2} - \sum |P_i|$ consisting of all the positive integers in the sequence

$$(i_1 - 1, i_2 - i_1 - 1, i_3 - i_2 - 1, \dots, i_\ell - i_{\ell-1} - 1, n - i_\ell).$$

Example: Let $n = 8$ and $D = \{P_1, \dots, P_8\}$ satisfying $(|P_1|, \dots, |P_8|) = (0, 1, 1, 2, 4, 4, 5, 6)$. Then D is a minimal staircase form. The set $\{i \mid |P_i| = i - 1\}$ equals $\{1, 2, 5\}$. The positive integers in the sequence $(1 - 1, 2 - 1 - 1, 5 - 2 - 1, 8 - 5)$ are $(2, 3)$, so the partition type of D is $(2, 3)$.



Let $p(k)$ denote the number of partitions of an integer k and Π_k denote the set of partitions of k .

Theorem 1.2 Let k be any positive integer. There are positive constants $c_1 = 8k + 5, c_2 = 2k + 1$ such that the following holds:

For integers n, d_1, d_2 satisfying $n \geq c_1, d_1 \geq c_2 n, d_2 \geq c_2 n$ and $d_1 + d_2 = \binom{n}{2} - k$, the vector space $(M_n)_{d_1, d_2}$ has dimension $p(k)$, and the $p(k)$ elements

$$\{ \text{a minimal staircase form of bi-degree } (d_1, d_2) \text{ and of partition type } \mu \}_{\mu \in \Pi_k}$$

form a basis of $(M_n)_{d_1, d_2}$.

Note that N.Bergeron and Z.Chen have found explicit bases for $(M_n)_{d_1, d_2}$ for certain bi-degrees using a different method [1].

1.2.2 For arbitrary k and n

Denote by $p(k)$ the partition number of k and by convention $p(0) = 1$ and $p(k) = 0$ for $k < 0$. Denote by $p(b, k)$ the partition number of k into no more than b parts, and by convention $p(0, k) = 0$ for $k > 0$, $p(b, 0) = 1$ for $b \geq 0$. One of our main results is as follows.

Theorem 1.3 *Let d_1, d_2 be non-negative integers d_1, d_2 with $d_1 + d_2 \leq \binom{n}{2}$. Define $k = \binom{n}{2} - d_1 - d_2$ and $\delta = \min(d_1, d_2)$. Then the coefficient of $q^{d_1}t^{d_2}$ in $C_n(q, t)$ is less than or equal to $p(\delta, k)$, and the equality holds if and only if one the following conditions holds:*

- $k \leq n - 3$, or
- $k = n - 2$ and $\delta = 1$, or
- $\delta = 0$.

This theorem is a consequence of Theorem C. It contains [8, Theorem 6] and a result of N.Bergeron and Z.Chen [1, Corollary 8.3.1] as special cases. In fact it proves [8, Conjecture 8]. We feel that the coefficient of $q^{d_1}t^{d_2}$ for general k can also be expressed in terms of partition numbers, only that the expression might be complicated. For example, we give the following conjecture which is verified for $6 \leq n \leq 10$.

Conjecture. Let n, d_1, d_2, δ, k be as in Theorem 1.3. If $n - 2 \leq k \leq 2n - 8$ and $\delta \geq k$, then the coefficient of $q^{d_1}t^{d_2}$ in $C_n(q, t)$ equals

$$p(k) - 2[p(0) + p(1) + \dots + p(k - n + 1)] - p(k - n + 2).$$

As a corollary of Theorem 1.3 , we can compute some higher degree terms of the specialization at $t = q$.

Corollary 1.4

$$C_n(q, q) = \sum_{k=0}^{n-3} \left(p(k) \left(\binom{n}{2} - 3k + 1 \right) + 2 \sum_{i=1}^{k-1} p(i, k) \right) q^{\binom{n}{2}-k} + (\text{lower degree terms}).$$

The following theorem immediately implies Theorem 1.3.

Theorem 1.5 *Let d_1, d_2 be non-negative integers d_1, d_2 with $d_1 + d_2 \leq \binom{n}{2}$. Define $k = \binom{n}{2} - d_1 - d_2$ and $\delta = \min(d_1, d_2)$. Then $\dim(M_n)_{d_1, d_2} \leq p(\delta, k)$, and the equality holds if and only if one the following conditions holds:*

- $k \leq n - 3$, or
- $k = n - 2$ and $\delta = 1$, or
- $\delta = 0$.

In case the equality holds, there is an explicit construction of a basis of $(M_n)_{d_1, d_2}$.

The idea of the construction of the basis in the above theorem consists of two parts:

(1) Prove that

$$\dim(M_n)_{d_1, d_2} \leq p(\delta, k)$$

using a new characterization of q, t -Catalan numbers. The characterization is as follows, and is discovered independently by A. Woo [10].

Let $\mathfrak{D}_n^{\text{catalan}}$ be the set consisting of $D \subset \mathbb{N} \times \mathbb{N}$, where D contains n points satisfying the following conditions.

- (a) If $(p, 0) \in D$ then $(i, 0) \in D, \forall i \in [0, p]$.
- (b) For any $p \in \mathbb{N}$,

$$\#\{j \mid (p + 1, j) \in D\} + \#\{j \mid (p, j) \in D\} \geq \max\{j \mid (p, j) \in D\} + 1.$$

(If $\{j \mid (p, j) \in D\} = \emptyset$, then we require that no point $(i, j) \in D$ satisfies $i \geq p$.) Denote by $\deg_x D$ (resp. $\deg_y D$) the sum of the first (resp. second) components of the n points in D .

Proposition 1.6 *The coefficient of $q^{d_1}t^{d_2}$ in the q, t -Catalan number $C_n(q, t)$ is equal to*

$$\#\{D \in \mathfrak{D}_n^{\text{catalan}} \mid \deg_x D = d_1, \deg_y D = d_2\}.$$

(2) Construct a set of $p(\delta, k)$ linearly independent elements in $(M_n)_{d_1, d_2}$. It seems difficult (as least to the authors) to test directly whether a given set of elements in $(M_n)_{d_1, d_2}$ are linearly independent. We define a map φ sending an alternating polynomial $f \in \mathbb{C}[\mathbf{x}, \mathbf{y}]^\epsilon$ to a polynomial ring

$$\mathbb{C}[\rho] := \mathbb{C}[\rho_1, \rho_2, \rho_3, \dots].$$

The map has two desirable properties: (i) for many f , $\varphi(f)$ can be easily computed, and (ii) for each bi-degree (d_1, d_2) , φ induces a morphism $\bar{\varphi} : (M_n)_{d_1, d_2} \rightarrow \mathbb{C}[\rho]$ of \mathbb{C} -modules. Then we use the fact the linear dependency is easier to check in $\mathbb{C}[\rho]$ than in $(M_n)_{d_1, d_2}$. The map φ is defined as below.

1.2.3 Maps φ and $\bar{\varphi}$.

(a) Define the map $\varphi : \mathfrak{D}_n \rightarrow \mathbb{Z}[\rho]$ as follows. Let $D = \{(a_1, b_1), \dots, (a_n, b_n)\} \in \mathfrak{D}_n, k = \binom{n}{2} - \sum_{i=1}^n (a_i + b_i)$, and define

$$\varphi(D) := (-1)^k \sum_{\sigma \in S_n} \text{sgn}(\sigma) \prod_{i=1}^n \left(\sum \rho_{w_1} \rho_{w_2} \cdots \rho_{w_{b_i}} \right),$$

where (w_1, \dots, w_{b_i}) in the sum $\sum \rho_{w_1} \rho_{w_2} \cdots \rho_{w_{b_i}}$ runs through the set

$$\{(w_1, \dots, w_{b_i}) \in \mathbb{N}^{b_i} \mid w_1 + \dots + w_{b_i} = \sigma(i) - 1 - a_i - b_i\},$$

with the convention that

$$\sum \rho_{w_1} \cdots \rho_{w_{b_i}} = \begin{cases} 0 & \text{if } \sigma(i) - 1 - a_i - b_i < 0; \\ 0 & \text{if } b_i = 0 \text{ and } \sigma(i) - 1 - a_i - b_i > 0; \\ 1 & \text{if } b_i = 0 \text{ and } \sigma(i) - 1 - a_i - b_i = 0. \end{cases}$$

(b) Here is an equivalent definition of $\varphi(D)$. Define the weight of ρ_i to be i for $i \in \mathbb{N}^+$ and define the weight of $\rho_0 = 1$ to be 0. Naturally the weight of any monomial $c\rho_{i_1}\dots\rho_{i_n}$ ($c \in \mathbb{Z}$) is defined to be $i_1 + \dots + i_n$. For $w \in \mathbb{N}$ and a power series $f \in \mathbb{Z}[[\rho_1, \rho_2, \dots]]$, denote by $\{f\}_w$ the sum of terms of weight- w in f , which is a polynomial. Define

$$h(b, w) := \{(1 + \rho_1 + \rho_2 + \dots)^b\}_w, \quad b \in \mathbb{N}, w \in \mathbb{Z}.$$

Naturally $h(b, w) = 0$ if $w < 0$. Also assume $(1 + \rho_1 + \rho_2 + \dots)^0 = 1$. Then

$$\varphi(D) = (-1)^k \begin{vmatrix} h(b_1, -|P_1|) & h(b_1, 1 - |P_1|) & h(b_1, 2 - |P_1|) & \dots & h(b_1, n - 1 - |P_1|) \\ h(b_2, -|P_2|) & h(b_2, 1 - |P_2|) & h(b_2, 2 - |P_2|) & \dots & h(b_2, n - 1 - |P_2|) \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ h(b_n, -|P_n|) & h(b_n, 1 - |P_n|) & h(b_n, 2 - |P_n|) & \dots & h(b_n, n - 1 - |P_n|) \end{vmatrix}.$$

(c) Let $D_1, \dots, D_\ell \in D'$ be of the same bi-degree and $\sum_{i=1}^\ell c_i D_i$ be the formal sum for any $c_i \in \mathbb{C}$ ($1 \leq i \leq \ell$). Define

$$\varphi\left(\sum_{i=1}^\ell c_i D_i\right) := \sum_{i=1}^\ell c_i \varphi(D_i).$$

For any bi-homogeneous alternating polynomials $f = \sum_{i=1}^\ell c_i \Delta(D_i) \in \mathbb{C}[\mathbf{x}, \mathbf{y}]^\epsilon$, we define

$$\varphi(f) := \varphi\left(\sum_{i=1}^\ell c_i D_i\right) = \sum_{i=1}^\ell c_i \varphi(D_i)$$

by abuse of notation. □

Proposition 1.7 Fix any pair of nonnegative integers (d_1, d_2) , the map φ induces a well-defined linear map

$$\bar{\varphi} : (M_n)_{d_1, d_2} \longrightarrow \mathbb{C}[\rho].$$

Moreover, this map $\bar{\varphi}$ is conjecturally injective. And our future work is to generalizing it to the case $I_n^m / (\mathbf{x}, \mathbf{y}) I_n^m$ for any positive integer m .

Acknowledgements

We are grateful to François Bergeron, Mahir Can, Jim Haglund, Nick Loehr, Alex Woo and Alex Yong for valuable discussions and correspondence.

References

- [1] N. Bergeron and Z. Chen, Basis of Diagonally Alternating Harmonic Polynomials for low degree, arXiv: 0905.0377.
- [2] A. M. Garsia and J. Haglund, A positivity result in the theory of Macdonald polynomials, Proc. Natl. Acad. Sci. USA **98** (2001), no. 8, 4313–4316 (electronic).

- [3] A. M. Garsia and J. Haglund, A proof of the q, t -Catalan positivity conjecture. LaCIM 2000 Conference on Combinatorics, Computer Science and Applications (Montreal, QC). *Discrete Math.* 256 (2002), no. 3, 677–717.
- [4] A. M. Garsia and M. Haiman, A Remarkable $q; t$ -Catalan sequence and q -Lagrange inversion, *J. Algebraic Combin.* 5 (1996), 191–244.
- [5] M. Haiman, Hilbert schemes, polygraphs and the Macdonald positivity conjecture, *J. Amer. Math. Soc.* 14 (2001), no. 4, 941–1006.
- [6] M. Haiman, Vanishing theorems and character formulas for the Hilbert scheme of points in the plane, *Invent. Math.* 149 (2002), no. 2, 371–407.
- [7] M. Haiman, Commutative algebra of n points in the plane, With an appendix by Ezra Miller. *Math. Sci. Res. Inst. Publ.*, 51, Trends in commutative algebra, 153–180, Cambridge Univ. Press, Cambridge, 2004.
- [8] K. Lee, L. Li, Notes on a minimal set of generators for the radical ideal defining the diagonal locus of $(\mathbb{C}^2)^n$, Arxiv math 0901.1176.
- [9] K. Lee, L. Li, q, t -Catalan numbers and generators for the radical ideal defining the diagonal locus of $(\mathbb{C}^2)^n$, ArXiv math 0909.1612.
- [10] A. Woo, private communication.