

Tiling groups and algorithms for tiling

Claire Kenyon

LRI, Bât 490, Université Paris-Sud, 91405 Orsay Cedex, France

We present algorithms for tiling a simply connected region of the plane (*i.e.* polygonal region without holes) with various kinds of tiles. Many of the algorithms use the concept of tiling group of J. H. Conway and Lagarias [1], ideas of W. P. Thurston on using the geometry of the group [12]. The concept of tiling groups is best illustrated by examples.

The first problem studied using tiling groups is the domino tiling problem. Given a simply connected polyomino P (polygon in \mathbf{R}^2 , with vertices in \mathbf{Z}^2 , horizontal or vertical edges, and no “holes”), find a tiling of P using dominoes, *i.e.* 1×2 and 2×1 rectangles, if it exists.

A moment's thought shows that the problem can be reduced *via* duality to perfect matching. However in [1] Conway and Lagarias used a radically different approach by introducing the *domino tiling group*. Let a represent a horizontal step to the right, a^{-1} a horizontal step to the left, b a vertical step going up, and b^{-1} a vertical step going down. Then the perimeters of the dominoes, read in a counterclockwise order from the lower left corner, correspond to the words $a^2ba^{-2}b^{-1}$ and $ab^2a^{-1}b^{-2}$. The perimeter of any simply connected polyomino P corresponds to a word $w(P)$.

Let $F = \langle a, b \rangle$ denote the free group on two generators and in general let $\langle A | R \rangle$ denote the group generated by the elements of A with relations $r = e$ for $r \in R$. (Thus the discrete plane \mathbf{Z}^2 has the group presentation $\mathbf{Z}^2 = \langle a, b | aba^{-1}b^{-1} \rangle$). The domino tiling group is defined as $G = \langle a, b | a^2ba^{-2}b^{-1}, ab^2a^{-1}b^{-2} \rangle$.

It is easy to see that if P is tileable by dominos, then $w(P)$ is equal to the identity in G [1]. Moreover, by studying the geometry of the Cayley graph of G , Thurston obtained an efficient and elegant algorithm for deciding if P is tileable and constructing a tiling if one exists [12].

The same approach can be used for other tiling problems.

- Tiling with “bars”, $1 \times m$ and $m \times 1$ rectangles with m fixed [2].
- Tiling with $m \times n$ and $n \times m$ rectangles with n, m fixed [2].
- When the tiling group is equal to \mathbf{Z}^2 , it suggests that the corresponding tiling problem may be “easy”. This is confirmed in the case of tiling with “bibones” [3].
- When the underlying planar lattice is the triangular lattice, one can also use tiling groups to get tiling algorithms for “leaning dominoes” and other families of tiles [10].
- One can also use the tiling group to obtain some information on tiling with “tetrominos”, polyominos of area 4 [9, 8], on tiling with “tribones” [12], and on the tileability of a planar polygon with squares [7].
- The tiling group approach can be used to give a new proof of the result that if a rectangle R is tiled with rectangles, each having at least one side of integral length, then R has a side of integral length [5].

- A generalization of the notion of tiling group can also be used to construct an example of a tiling problem which is not “Escher-equivalent” to any problem in which all the tiles are polygonal [4].

References

- [1] J. H. Conway, J. C. Lagarias, “Tiling with polyominoes and combinatorial group theory”, *J. Comb. Theory, Ser. A* 53 (1990), 183-208.
- [2] Claire Kenyon and Richard Kenyon, “Tiling a polygon with rectangles”, 33rd Annual Symposium on Foundations of Computer Science (FOCS), Pittsburgh, PA, 1992, 610-619.
- [3] Claire Kenyon and Eric Rémila, “Perfect matchings on the triangular lattice”, *Discrete Mathematics*, 152 (1996), 191-210.
- [4] R. Kenyon, A group of paths in the plane, *Trans. AMS* 348,(1996):3155-3172.
- [5] R. Kenyon, A note on tiling with integer-sided rectangles, *J. Combin. Thy A* 74, No. 2 (1996):321-332.
- [6] R. Kenyon, Tilings of convex polygons, *Ann. Inst. Fourier*, 47, 3(1997),929-944.
- [7] R. Kenyon, “Tiling with squares and square-tileable surfaces”, preprint, 1998.
- [8] R. Muchnik and I. Pak, “On tilings by ribbon tetrominoes”, Yale university preprint, 1998.
- [9] J. Propp, “A Pedestrian Approach to a Method of Conway, or, A Tale of Two Cities “, *Mathematics Magazine* 70,December 1997, 327-340.
- [10] E. Rémila, “Tiling a figure using a height in a tree”, *Proc. of the 7th Ann. ACM-SIAM Symp. on Disc. Algorithms*, 1996, pp 168-174.
- [11] E. Rémila, “Tiling groups: new applications in the triangular lattice”, *Discrete and Computational Geometry* 1998 20, 189-204.
- [12] William P. Thurston, “Conway’s Tiling Groups”, *Amer. Math. Monthly*, 97, Oct. 1990, pp. 757-773.